

#17 IS 09/688,851



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, 15/82, 15/65, G01N 33/52		A2	(11) International Publication Number: WO 97/41228
			(43) International Publication Date: 6 November 1997 (06.11.97)
(21) International Application Number: PCT/US97/07688		<p>Circle, Johnston, IA 50131 (US). SCELONGE, Christopher [US/US]; 5314 Ovid Avenue, Des Moines, IA 50310 (US). MILLER, Mike [US/US]; 59 W. Buchanan Street, Winterset, IA 50273 (US). SANDAHL, Gary [US/US]; 1649 Thornwood Road, West Des Moines, IA 50265 (US). WANG, Lijuan [CN/US]; 4728 80th Place, Urbandale, IA 50322 (US).</p> <p>(74) Agents: BENT, Stephen, A. et al.; Foley & Lardner, Suite 500, 3000 K Street, N.W., Washington, DC 20007-5109 (US).</p> <p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU. ARIPO patent (GH, KE, LS, MW, SD, SZ, UG). Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM). European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published Without international search report and to be republished upon receipt of that report.</p>	
(22) International Filing Date: 1 May 1997 (01.05.97)			
(30) Priority Data: 60/016,345 1 May 1996 (01.05.96) US			
(60) Parent Application or Grant (63) Related by Continuation US 60/016,345 (CIP) Filed on 1 May 1996 (01.05.96)			
(71) Applicant (for all designated States except US): PIONEER HIBRED INTERNATIONAL, INC. [US/US]; Darwin Building, 7100 N.W. 62nd Avenue, P.O. Box 1000, Johnston, IA 50131 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): GORDON-KAMM, William [US/US]; 3916 67th Street, Urbandale, IA 50322 (US). PIERCE, Dorothy, A. [US/US]; 9124 Hammontree Drive, Urbandale, IA 50322 (US). BOWEN, Benjamin [GB/US]; 3008 36th Street, Des Moines, IA 50310 (US). BIDNEY, Dennis [US/US]; 8385 Plum Drive, Urbandale, IA 50322 (US). ROSS, Margit [US/US]; 5534 Kensington			
(54) Title: USE OF THE GREEN FLUORESCENT PROTEIN AS A SCREENABLE MARKER FOR PLANT TRANSFORMATION			
(57) Abstract			
<p>A method for the production of transgenic plants is provided in which a vector carrying a gene encoding the green fluorescent protein is introduced into cells, the cells are screened for the protein and transformed cells are selected and regenerated. The cellular toxicity of the green fluorescent protein is circumvented by regulating expression of the gene encoding the protein or directing the protein to a subcellular compartment where it is not toxic to the cell. DNA constructs are provided for cell transformation in which the expression of a gene encoding the green fluorescent protein is placed under the control of an inducible promoter. In addition, DNA constructs are provided in which a nucleotide sequence encoding the green fluorescent protein is operably linked to a signal sequence which directs the expressed protein to a subcellular compartment where the protein is not toxic to the cell. Oxidative stress to plant cells transformed with GFP also can be ameliorated by transforming cells with an expression vector comprising genes encoding GFP and an oxygen scavenger enzyme such as superoxide dismutase. The toxicity of GFP in transformed plants can be eliminated by excising the screenable marker gene following detection of transformed cells or sectors. The FLP/FRT system is used in conjunction with GFP as a visible marker for transformation and FRT excision. A nucleotide sequence optimized for expression of the green fluorescent protein in plants is also provided.</p>			
<p>Sequence listing (continued) (Table 1) (Table 2) (Table 3) (Table 4) (Table 5) (Table 6) (Table 7) (Table 8) (Table 9) (Table 10) (Table 11) (Table 12) (Table 13) (Table 14) (Table 15) (Table 16) (Table 17) (Table 18) (Table 19) (Table 20) (Table 21) (Table 22) (Table 23) (Table 24) (Table 25) (Table 26) (Table 27) (Table 28) (Table 29) (Table 30) (Table 31) (Table 32) (Table 33) (Table 34) (Table 35) (Table 36) (Table 37) (Table 38) (Table 39) (Table 40) (Table 41) (Table 42) (Table 43) (Table 44) (Table 45) (Table 46) (Table 47) (Table 48) (Table 49) (Table 50) (Table 51) (Table 52) (Table 53) (Table 54) (Table 55) (Table 56) (Table 57) (Table 58) (Table 59) (Table 60) (Table 61) (Table 62) (Table 63) (Table 64) (Table 65) (Table 66) (Table 67) (Table 68) (Table 69) (Table 70) (Table 71) (Table 72) (Table 73) (Table 74) (Table 75) (Table 76) (Table 77) (Table 78) (Table 79) (Table 80) (Table 81) (Table 82) (Table 83) (Table 84) (Table 85) (Table 86) (Table 87) (Table 88) (Table 89) (Table 90) (Table 91) (Table 92) (Table 93) (Table 94) (Table 95) (Table 96) (Table 97) (Table 98) (Table 99) (Table 100) (Table 101) (Table 102) (Table 103) (Table 104) (Table 105) (Table 106) (Table 107) (Table 108) (Table 109) (Table 110) (Table 111) (Table 112) (Table 113) (Table 114) (Table 115) (Table 116) (Table 117) (Table 118) (Table 119) (Table 120) (Table 121) (Table 122) (Table 123) (Table 124) (Table 125) (Table 126) (Table 127) (Table 128) (Table 129) (Table 130) (Table 131) (Table 132) (Table 133) (Table 134) (Table 135) (Table 136) (Table 137) (Table 138) (Table 139) (Table 140) (Table 141) (Table 142) (Table 143) (Table 144) (Table 145) (Table 146) (Table 147) (Table 148) (Table 149) (Table 150) (Table 151) (Table 152) (Table 153) (Table 154) (Table 155) (Table 156) (Table 157) (Table 158) (Table 159) (Table 160) (Table 161) (Table 162) (Table 163) (Table 164) (Table 165) (Table 166) (Table 167) (Table 168) (Table 169) (Table 170) (Table 171) (Table 172) (Table 173) (Table 174) (Table 175) (Table 176) (Table 177) (Table 178) (Table 179) (Table 180) (Table 181) (Table 182) (Table 183) (Table 184) (Table 185) (Table 186) (Table 187) (Table 188) (Table 189) (Table 190) (Table 191) (Table 192) (Table 193) (Table 194) (Table 195) (Table 196) (Table 197) (Table 198) (Table 199) (Table 200) (Table 201) (Table 202) (Table 203) (Table 204) (Table 205) (Table 206) (Table 207) (Table 208) (Table 209) (Table 210) (Table 211) (Table 212) (Table 213) (Table 214) (Table 215) (Table 216) (Table 217) (Table 218) (Table 219) (Table 220) (Table 221) (Table 222) (Table 223) (Table 224) (Table 225) (Table 226) (Table 227) (Table 228) (Table 229) (Table 230) (Table 231) (Table 232) (Table 233) (Table 234) (Table 235) (Table 236) (Table 237) (Table 238) (Table 239) (Table 240) (Table 241) (Table 242) (Table 243) (Table 244) (Table 245) (Table 246) (Table 247) (Table 248) (Table 249) (Table 250) (Table 251) (Table 252) (Table 253) (Table 254) (Table 255) (Table 256) (Table 257) (Table 258) (Table 259) (Table 260) (Table 261) (Table 262) (Table 263) (Table 264) (Table 265) (Table 266) (Table 267) (Table 268) (Table 269) (Table 270) (Table 271) (Table 272) (Table 273) (Table 274) (Table 275) (Table 276) (Table 277) (Table 278) (Table 279) (Table 280) (Table 281) (Table 282) (Table 283) (Table 284) (Table 285) (Table 286) (Table 287) (Table 288) (Table 289) (Table 290) (Table 291) (Table 292) (Table 293) (Table 294) (Table 295) (Table 296) (Table 297) (Table 298) (Table 299) (Table 300) (Table 301) (Table 302) (Table 303) (Table 304) (Table 305) (Table 306) (Table 307) (Table 308) (Table 309) (Table 310) (Table 311) (Table 312) (Table 313) (Table 314) (Table 315) (Table 316) (Table 317) (Table 318) (Table 319) (Table 320) (Table 321) (Table 322) (Table 323) (Table 324) (Table 325) (Table 326) (Table 327) (Table 328) (Table 329) (Table 330) (Table 331) (Table 332) (Table 333) (Table 334) (Table 335) (Table 336) (Table 337) (Table 338) (Table 339) (Table 340) (Table 341) (Table 342) (Table 343) (Table 344) (Table 345) (Table 346) (Table 347) (Table 348) (Table 349) (Table 350) (Table 351) (Table 352) (Table 353) (Table 354) (Table 355) (Table 356) (Table 357) (Table 358) (Table 359) (Table 360) (Table 361) (Table 362) (Table 363) (Table 364) (Table 365) (Table 366) (Table 367) (Table 368) (Table 369) (Table 370) (Table 371) (Table 372) (Table 373) (Table 374) (Table 375) (Table 376) (Table 377) (Table 378) (Table 379) (Table 380) (Table 381) (Table 382) (Table 383) (Table 384) (Table 385) (Table 386) (Table 387) (Table 388) (Table 389) (Table 390) (Table 391) (Table 392) (Table 393) (Table 394) (Table 395) (Table 396) (Table 397) (Table 398) (Table 399) (Table 400) (Table 401) (Table 402) (Table 403) (Table 404) (Table 405) (Table 406) (Table 407) (Table 408) (Table 409) (Table 410) (Table 411) (Table 412) (Table 413) (Table 414) (Table 415) (Table 416) (Table 417) (Table 418) (Table 419) (Table 420) (Table 421) (Table 422) (Table 423) (Table 424) (Table 425) (Table 426) (Table 427) (Table 428) (Table 429) (Table 430) (Table 431) (Table 432) (Table 433) (Table 434) (Table 435) (Table 436) (Table 437) (Table 438) (Table 439) (Table 440) (Table 441) (Table 442) (Table 443) (Table 444) (Table 445) (Table 446) (Table 447) (Table 448) (Table 449) (Table 450) (Table 451) (Table 452) (Table 453) (Table 454) (Table 455) (Table 456) (Table 457) (Table 458) (Table 459) (Table 460) (Table 461) (Table 462) (Table 463) (Table 464) (Table 465) (Table 466) (Table 467) (Table 468) (Table 469) (Table 470) (Table 471) (Table 472) (Table 473) (Table 474) (Table 475) (Table 476) (Table 477) (Table 478) (Table 479) (Table 480) (Table 481) (Table 482) (Table 483) (Table 484) (Table 485) (Table 486) (Table 487) (Table 488) (Table 489) (Table 490) (Table 491) (Table 492) (Table 493) (Table 494) (Table 495) (Table 496) (Table 497) (Table 498) (Table 499) (Table 500) (Table 501) (Table 502) (Table 503) (Table 504) (Table 505) (Table 506) (Table 507) (Table 508) (Table 509) (Table 510) (Table 511) (Table 512) (Table 513) (Table 514) (Table 515) (Table 516) (Table 517) (Table 518) (Table 519) (Table 520) (Table 521) (Table 522) (Table 523) (Table 524) (Table 525) (Table 526) (Table 527) (Table 528) (Table 529) (Table 530) (Table 531) (Table 532) (Table 533) (Table 534) (Table 535) (Table 536) (Table 537) (Table 538) (Table 539) (Table 540) (Table 541) (Table 542) (Table 543) (Table 544) (Table 545) (Table 546) (Table 547) (Table 548) (Table 549) (Table 550) (Table 551) (Table 552) (Table 553) (Table 554) (Table 555) (Table 556) (Table 557) (Table 558) (Table 559) (Table 560) (Table 561) (Table 562) (Table 563) (Table 564) (Table 565) (Table 566) (Table 567) (Table 568) (Table 569) (Table 570) (Table 571) (Table 572) (Table 573) (Table 574) (Table 575) (Table 576) (Table 577) (Table 578) (Table 579) (Table 580) (Table 581) (Table 582) (Table 583) (Table 584) (Table 585) (Table 586) (Table 587) (Table 588) (Table 589) (Table 590) (Table 591) (Table 592) (Table 593) (Table 594) (Table 595) (Table 596) (Table 597) (Table 598) (Table 599) (Table 600) (Table 601) (Table 602) (Table 603) (Table 604) (Table 605) (Table 606) (Table 607) (Table 608) (Table 609) (Table 610) (Table 611) (Table 612) (Table 613) (Table 614) (Table 615) (Table 616) (Table 617) (Table 618) (Table 619) (Table 620) (Table 621) (Table 622) (Table 623) (Table 624) (Table 625) (Table 626) (Table 627) (Table 628) (Table 629) (Table 630) (Table 631) (Table 632) (Table 633) (Table 634) (Table 635) (Table 636) (Table 637) (Table 638) (Table 639) (Table 640) (Table 641) (Table 642) (Table 643) (Table 644) (Table 645) (Table 646) (Table 647) (Table 648) (Table 649) (Table 650) (Table 651) (Table 652) (Table 653) (Table 654) (Table 655) (Table 656) (Table 657) (Table 658) (Table 659) (Table 660) (Table 661) (Table 662) (Table 663) (Table 664) (Table 665) (Table 666) (Table 667) (Table 668) (Table 669) (Table 670) (Table 671) (Table 672) (Table 673) (Table 674) (Table 675) (Table 676) (Table 677) (Table 678) (Table 679) (Table 680) (Table 681) (Table 682) (Table 683) (Table 684) (Table 685) (Table 686) (Table 687) (Table 688) (Table 689) (Table 690) (Table 691) (Table 692) (Table 693) (Table 694) (Table 695) (Table 696) (Table 697) (Table 698) (Table 699) (Table 700) (Table 701) (Table 702) (Table 703) (Table 704) (Table 705) (Table 706) (Table 707) (Table 708) (Table 709) (Table 710) (Table 711) (Table 712) (Table 713) (Table 714) (Table 715) (Table 716) (Table 717) (Table 718) (Table 719) (Table 720) (Table 721) (Table 722) (Table 723) (Table 724) (Table 725) (Table 726) (Table 727) (Table 728) (Table 729) (Table 730) (Table 731) (Table 732) (Table 733) (Table 734) (Table 735) (Table 736) (Table 737) (Table 738) (Table 739) (Table 740) (Table 741) (Table 742) (Table 743) (Table 744) (Table 745) (Table 746) (Table 747) (Table 748) (Table 749) (Table 750) (Table 751) (Table 752) (Table 753) (Table 754) (Table 755) (Table 756) (Table 757) (Table 758) (Table 759) (Table 760) (Table 761) (Table 762) (Table 763) (Table 764) (Table 765) (Table 766) (Table 767) (Table 768) (Table 769) (Table 770) (Table 771) (Table 772) (Table 773) (Table 774) (Table 775) (Table 776) (Table 777) (Table 778) (Table 779) (Table 780) (Table 781) (Table 782) (Table 783) (Table 784) (Table 785) (Table 786) (Table 787) (Table 788) (Table 789) (Table 790) (Table 791) (Table 792) (Table 793) (Table 794) (Table 795) (Table 796) (Table 797) (Table 798) (Table 799) (Table 800) (Table 801) (Table 802) (Table 803) (Table 804) (Table 805) (Table 806) (Table 807) (Table 808) (Table 809) (Table 810) (Table 811) (Table 812) (Table 813) (Table 814) (Table 815) (Table 816) (Table 817) (Table 818) (Table 819) (Table 820) (Table 821) (Table 822) (Table 823) (Table 824) (Table 825) (Table 826) (Table 827) (Table 828) (Table 829) (Table 830) (Table 831) (Table 832) (Table 833) (Table 834) (Table 835) (Table 836) (Table 837) (Table 838) (Table 839) (Table 840) (Table 841) (Table 842) (Table 843) (Table 844) (Table 845) (Table 846) (Table 847) (Table 848) (Table 849) (Table 850) (Table 851) (Table 852) (Table 853) (Table 854) (Table 855) (Table 856) (Table 857) (Table 858) (Table 859) (Table 860) (Table 861) (Table 862) (Table 863) (Table 864) (Table 865) (Table 866) (Table 867) (Table 868) (Table 869) (Table 870) (Table 871) (Table 872) (Table 873) (Table 874) (Table 875) (Table 876) (Table 877) (Table 878) (Table 879) (Table 880) (Table 881) (Table 882) (Table 883) (Table 884) (Table 885) (Table 886) (Table 887) (Table 888) (Table 889) (Table 890) (Table 891) (Table 892) (Table 893) (Table 894) (Table 895) (Table 896) (Table 897) (Table 898) (Table 899) (Table 900) (Table 901) (Table 902) (Table 903) (Table 904) (Table 905) (Table 906) (Table 907) (Table 908) (Table 909) (Table 910) (Table 911) (Table 912) (Table 913) (Table 914) (Table 915) (Table 916) (Table 917) (Table 918) (Table 919) (Table 920) (Table 921) (Table 922) (Table 923) (Table 924) (Table 925) (Table 926) (Table 927) (Table 928) (Table 929) (Table 930) (Table 931) (Table 932) (Table 933) (Table 934) (Table 935) (Table 936) (Table 937) (Table 938) (Table 939) (Table 940) (Table 941) (Table 942) (Table 943) (Table 944) (Table 945) (Table 946) (Table 947) (Table 948) (Table 949) (Table 950) (Table 951) (Table 952) (Table 953) (Table 954) (Table 955) (Table 956) (Table 957) (Table 958) (Table 959) (Table 960) (Table 961) (Table 962) (Table 963) (Table 964) (Table 965) (Table 966) (Table 967) (Table 968) (Table 969) (Table 970) (Table 971) (Table 972) (Table 973) (Table 974) (Table 975) (Table 976) (Table 977) (Table 978) (Table 979) (Table 980) (Table 981) (Table 982) (Table 983) (Table 984) (Table 985) (Table 986) (Table 987) (Table 988) (Table 989) (Table 990) (Table 991) (Table 992) (Table 993) (Table 994) (Table 995) (Table 996) (Table 997) (Table 998) (Table 999) (Table 1000) (Table 1001) (Table 1002) (Table 1003) (Table 1004) (Table 1005) (Table 1006) (Table 1007) (Table 1008) (Table 1009) (Table 1010) (Table 1011) (Table 1012) (Table 1013) (Table 1014) (Table 1015) (Table 1016) (Table 1017) (Table 1018) (Table 1019) (Table 1020) (Table 1021) (Table 1022) (Table 1023) (Table 1024) (Table 1025) (Table 1026) (Table 1027) (Table 1028) (Table 1029) (Table 1030) (Table 1031) (Table 1032) (Table 1033) (Table 1034) (Table 1035) (Table 1036) (Table 1037) (Table 1038) (Table 1039) (Table 1040) (Table 1041) (Table 1042) (Table 1043) (Table 1044) (Table 1045) (Table 1046) (Table 1047) (Table 1048) (Table 1049) (Table 1050) (Table 1051) (Table 1052) (Table 1053) (Table 1054) (Table 1055) (Table 1056) (Table 1057) (Table 1058) (Table 1059) (Table 1060) (Table 1061) (Table 1062) (Table 1063) (Table 1064) (Table 1065) (Table 1066) (Table 1067) (Table 1068) (Table 1069) (Table 1070) (Table 1071) (Table 1072) (Table 1073) (Table 1074) (Table 1075) (Table 1076) (Table 1077) (Table 1078) (Table 1079) (Table 1080) (Table 1081) (Table 1082) (Table 1083) (Table 1084) (Table 1085) (Table 1086) (Table 1087) (Table 1088) (Table 1089) (Table 1090) (Table 1091) (Table 1092) (Table 1093) (Table 1094) (Table 1095) (Table 1096) (Table 1097) (Table 1098) (Table 1099) (Table 1100) (Table 1101) (Table 1102) (Table 1103) (Table 1104) (Table 1105) (Table 1106) (Table 1107) (Table 1108) (Table 1109) (Table 1110) (Table 1111) (Table 1112) (Table 1113) (Table 1114) (Table 1115) (Table 1116) (Table 1117) (Table 1118) (Table 1119) (Table 1120) (Table 1121) (Table 1122) (Table 1123) (Table 1124) (Table 1125) (Table 1126) (Table 1127) (Table 1128) (Table 1129) (Table 1130) (Table 1131) (Table 1132) (Table 1133) (Table 1134) (Table 1135) (Table 1136) (Table 1137) (Table 1138) (Table 1139) (Table 1140) (Table 1141) (Table 1142) (Table 1143) (Table 1144) (Table 1145) (Table 1146) (Table 1147) (Table 1148) (Table 1149) (Table 1150) (Table 1151) (Table 1152) (Table 1153) (Table 1154) (Table 1155) (Table 1156) (Table 1157) (Table 1158) (Table 1159) (Table 1160) (Table 1161) (Table 1162) (Table 1163) (Table 1164) (Table 1165) (Table 1166) (Table 1167) (Table 1168) (Table 1169) (Table 1170) (Table 1171) (Table 1172) (Table 1173) (Table 1174) (Table 1175) (Table 1176) (Table 1177) (Table 1178) (Table 1179) (Table 1180) (Table 1181) (Table 1182) (Table 1183) (Table 1184) (Table 1185) (Table 1186) (Table 1187) (Table 1188) (Table 1189) (Table 1190) (Table 1191) (Table 1192) (Table 1193) (Table 1194) (Table 1195) (Table 1196) (Table 1197) (Table 1198) (Table 1199) (Table 1200) (Table 1201) (Table 1202) (Table 1203) (Table 1204) (Table 1205) (Table 1206) (Table 1207) (Table 1208) (Table 1209) (Table 1210) (Table 1211) (Table 1212) (Table 1213) (Table 1214) (Table 1215) (Table 1216) (Table 1217) (Table 1218) (Table 1219) (Table 1220) (Table 1221) (Table 1222) (Table 1223) (Table 1224) (Table 1225) (Table 1226) (Table 1227) (Table 1228) (Table 1229) (Table 1230) (Table 1231) (Table 1232) (Table 1233) (Table 1234) (Table 1235) (Table 1236) (Table 1237) (Table 1238) (Table 1239) (Table 1240) (Table 1241) (Table 1242) (Table 1243) (Table 1244) (Table 1245) (Table 1246) (Table 1247) (Table 1248) (Table 1249) (Table 1250) (Table 1251) (Table 1252) (Table 1253) (Table 1254) (Table 1255) (Table 1256) (Table 1257) (Table 1258) (Table 1259) (Table 1260) (Table 1261) (Table 1262) (Table 1263) (Table 1264) (Table 1265) (Table 1266) (Table 1267) (Table 1268) (Table 1269) (Table 1270) (Table 1271) (Table 1272) (Table 1273) (Table 1274) (Table 1275) (Table 1276) (Table 1277) (Table 1278) (Table 1279) (Table 1280) (Table 1281) (Table 1282) (Table 1283) (Table 1284) (Table 1285) (Table 1286) (Table 1287) (Table 1288) (Table 1289) (Table 1290) (Table 1291) (Table 1292) (Table 1293) (Table 1294) (Table 1295) (Table 1296) (Table 1297) (Table 1298) (Table 1299) (Table 1300) (Table 1301) (Table 1302) (Table 1303) (Table 1304) (Table 1305) (Table 1306) (Table 1307) (Table 1308) (Table 1309) (Table 1310) (Table 1311) (Table 1312) (Table 1313) (Table 1314) (Table 1315) (Table 1316) (Table 1317) (Table 1318) (Table 1319) (Table 1320) (Table 1321) (Table 1322) (Table 1323) (Table 1324) (Table 1325) (Table 1326) (Table 1327) (Table 1328) (Table 1329) (Table 1330) (Table 1331) (Table 1332) (Table 1333) (Table 1334) (Table 1335) (Table 1336) (Table 1337) (Table 1338) (Table 1339) (Table 1340) (Table 1341) (Table 1342) (Table 1343) (Table 1344) (Table 1345) (Table 1346) (Table 1347) (Table 1348) (Table 1349) (Table 1350) (Table 1351) (Table 1352) (Table 1353) (Table 1354) (Table 1355) (Table 1356) (Table 1357) (Table 1358) (Table 1359) (Table 1360) (Table 1361) (Table 1362) (Table 1363) (Table 1364) (Table 1365) (Table 1366) (Table 1367) (Table 1368) (Table 1369) (Table 1370) (Table 1371) (Table 1372) (Table 1373) (Table 1374) (Table 1375) (Table 1376) (Table 1377) (Table 1378) (Table 1379) (Table 1380) (Table 1381) (Table 1382) (Table 1383) (Table 1384) (Table 1385) (Table 1386) (Table 1387) (Table 1388) (Table 1389) (Table 1390) (Table 1391) (Table 1392) (Table 1393) (Table 1394) (Table 1395) (Table 1396) (Table 1397) (Table 1398) (Table 1399) (Table 1400) (Table 1401) (Table 1402) (Table 1403) (Table 1404) (Table 1405) (Table 1406) (Table 1407) (Table 1408) (Table 1409) (Table 1410) (Table 1411) (Table 1412) (Table 1413) (Table 1414) (Table 1415) (Table 1416) (Table 1417) (Table 1418) (Table 1419) (Table 1420) (Table 1421) (Table 1422) (Table 1423) (Table 1424) (Table 1425) (Table 1426) (Table 1427) (Table 1428) (Table 1429) (Table 1430) (Table 1431) (Table 1432) (Table 1433) (Table 1434) (Table 1435) (Table 1436) (Table 1437) (Table 1438) (Table 1439) (Table 1440) (Table 1441) (Table 1442) (Table 1443) (Table 1444) (Table 1445) (Table 1446) (Table 1447) (Table 1448) (Table 1449) (Table 1450) (Table 1451) (Table 1452) (Table 1453) (Table 1454) (Table 1455) (Table 1456) (Table 1457) (Table 1458) (Table 1459) (Table 1460) (Table 1461) (Table 1462) (Table 1463) (Table 1464) (Table 1465) (Table 1466) (Table 1467) (Table 1468) (Table 1469) (Table 1470) (Table 1471) (Table 1472) (Table 1473) (Table 1474) (Table 1475) (Table 1476) (Table 1477) (</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

USE OF THE GREEN FLUORESCENT PROTEIN AS A SCREENABLE
MARKER FOR PLANT TRANSFORMATION

BACKGROUND OF THE INVENTION

1. Field of the Invention

5 The present invention relates to a method of plant transformation in which a DNA construct carrying a gene encoding the green fluorescent protein (GFP) is introduced into plant cells which are then screened for the presence of the protein and transformed cells are
10 regenerated into transgenic plants. In particular, the present invention provides methods for circumventing the cellular toxicity of the GFP by regulating expression of the gene encoding the protein or directing the protein to a subcellular compartment where it is not toxic to the
15 cell. The present invention provides DNA constructs for cell transformation in which expression of a gene encoding the GFP is placed under the control of an inducible, constitutive or tissue-specific promoter. In addition, DNA constructs are provided in which a
20 nucleotide sequence encoding the GFP is operably linked to a signal or targeting sequence which directs the expressed protein to a subcellular compartment where the protein is not toxic to the cell. Moreover, the present invention provides a nucleotide sequence encoding GFP
25 that is optimized for expression of the GFP gene in

plants and to GFP-encoding nucleotide sequences that code for light-shifted versions of GFP. The present invention also provides a method for selecting plant cells transformed with a gene encoding a screenable marker flanked on the 5-prime and 3-prime ends with a recombina-
5
10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995
1000
1005
1010
1015
1020
1025
1030
1035
1040
1045
1050
1055
1060
1065
1070
1075
1080
1085
1090
1095
1100
1105
1110
1115
1120
1125
1130
1135
1140
1145
1150
1155
1160
1165
1170
1175
1180
1185
1190
1195
1200
1205
1210
1215
1220
1225
1230
1235
1240
1245
1250
1255
1260
1265
1270
1275
1280
1285
1290
1295
1300
1305
1310
1315
1320
1325
1330
1335
1340
1345
1350
1355
1360
1365
1370
1375
1380
1385
1390
1395
1400
1405
1410
1415
1420
1425
1430
1435
1440
1445
1450
1455
1460
1465
1470
1475
1480
1485
1490
1495
1500
1505
1510
1515
1520
1525
1530
1535
1540
1545
1550
1555
1560
1565
1570
1575
1580
1585
1590
1595
1600
1605
1610
1615
1620
1625
1630
1635
1640
1645
1650
1655
1660
1665
1670
1675
1680
1685
1690
1695
1700
1705
1710
1715
1720
1725
1730
1735
1740
1745
1750
1755
1760
1765
1770
1775
1780
1785
1790
1795
1800
1805
1810
1815
1820
1825
1830
1835
1840
1845
1850
1855
1860
1865
1870
1875
1880
1885
1890
1895
1900
1905
1910
1915
1920
1925
1930
1935
1940
1945
1950
1955
1960
1965
1970
1975
1980
1985
1990
1995
2000
2005
2010
2015
2020
2025
2030
2035
2040
2045
2050
2055
2060
2065
2070
2075
2080
2085
2090
2095
2100
2105
2110
2115
2120
2125
2130
2135
2140
2145
2150
2155
2160
2165
2170
2175
2180
2185
2190
2195
2200
2205
2210
2215
2220
2225
2230
2235
2240
2245
2250
2255
2260
2265
2270
2275
2280
2285
2290
2295
2300
2305
2310
2315
2320
2325
2330
2335
2340
2345
2350
2355
2360
2365
2370
2375
2380
2385
2390
2395
2400
2405
2410
2415
2420
2425
2430
2435
2440
2445
2450
2455
2460
2465
2470
2475
2480
2485
2490
2495
2500
2505
2510
2515
2520
2525
2530
2535
2540
2545
2550
2555
2560
2565
2570
2575
2580
2585
2590
2595
2600
2605
2610
2615
2620
2625
2630
2635
2640
2645
2650
2655
2660
2665
2670
2675
2680
2685
2690
2695
2700
2705
2710
2715
2720
2725
2730
2735
2740
2745
2750
2755
2760
2765
2770
2775
2780
2785
2790
2795
2800
2805
2810
2815
2820
2825
2830
2835
2840
2845
2850
2855
2860
2865
2870
2875
2880
2885
2890
2895
2900
2905
2910
2915
2920
2925
2930
2935
2940
2945
2950
2955
2960
2965
2970
2975
2980
2985
2990
2995
3000
3005
3010
3015
3020
3025
3030
3035
3040
3045
3050
3055
3060
3065
3070
3075
3080
3085
3090
3095
3100
3105
3110
3115
3120
3125
3130
3135
3140
3145
3150
3155
3160
3165
3170
3175
3180
3185
3190
3195
3200
3205
3210
3215
3220
3225
3230
3235
3240
3245
3250
3255
3260
3265
3270
3275
3280
3285
3290
3295
3300
3305
3310
3315
3320
3325
3330
3335
3340
3345
3350
3355
3360
3365
3370
3375
3380
3385
3390
3395
3400
3405
3410
3415
3420
3425
3430
3435
3440
3445
3450
3455
3460
3465
3470
3475
3480
3485
3490
3495
3500
3505
3510
3515
3520
3525
3530
3535
3540
3545
3550
3555
3560
3565
3570
3575
3580
3585
3590
3595
3600
3605
3610
3615
3620
3625
3630
3635
3640
3645
3650
3655
3660
3665
3670
3675
3680
3685
3690
3695
3700
3705
3710
3715
3720
3725
3730
3735
3740
3745
3750
3755
3760
3765
3770
3775
3780
3785
3790
3795
3800
3805
3810
3815
3820
3825
3830
3835
3840
3845
3850
3855
3860
3865
3870
3875
3880
3885
3890
3895
3900
3905
3910
3915
3920
3925
3930
3935
3940
3945
3950
3955
3960
3965
3970
3975
3980
3985
3990
3995
4000
4005
4010
4015
4020
4025
4030
4035
4040
4045
4050
4055
4060
4065
4070
4075
4080
4085
4090
4095
4100
4105
4110
4115
4120
4125
4130
4135
4140
4145
4150
4155
4160
4165
4170
4175
4180
4185
4190
4195
4200
4205
4210
4215
4220
4225
4230
4235
4240
4245
4250
4255
4260
4265
4270
4275
4280
4285
4290
4295
4300
4305
4310
4315
4320
4325
4330
4335
4340
4345
4350
4355
4360
4365
4370
4375
4380
4385
4390
4395
4400
4405
4410
4415
4420
4425
4430
4435
4440
4445
4450
4455
4460
4465
4470
4475
4480
4485
4490
4495
4500
4505
4510
4515
4520
4525
4530
4535
4540
4545
4550
4555
4560
4565
4570
4575
4580
4585
4590
4595
4600
4605
4610
4615
4620
4625
4630
4635
4640
4645
4650
4655
4660
4665
4670
4675
4680
4685
4690
4695
4700
4705
4710
4715
4720
4725
4730
4735
4740
4745
4750
4755
4760
4765
4770
4775
4780
4785
4790
4795
4800
4805
4810
4815
4820
4825
4830
4835
4840
4845
4850
4855
4860
4865
4870
4875
4880
4885
4890
4895
4900
4905
4910
4915
4920
4925
4930
4935
4940
4945
4950
4955
4960
4965
4970
4975
4980
4985
4990
4995
5000
5005
5010
5015
5020
5025
5030
5035
5040
5045
5050
5055
5060
5065
5070
5075
5080
5085
5090
5095
5100
5105
5110
5115
5120
5125
5130
5135
5140
5145
5150
5155
5160
5165
5170
5175
5180
5185
5190
5195
5200
5205
5210
5215
5220
5225
5230
5235
5240
5245
5250
5255
5260
5265
5270
5275
5280
5285
5290
5295
5300
5305
5310
5315
5320
5325
5330
5335
5340
5345
5350
5355
5360
5365
5370
5375
5380
5385
5390
5395
5400
5405
5410
5415
5420
5425
5430
5435
5440
5445
5450
5455
5460
5465
5470
5475
5480
5485
5490
5495
5500
5505
5510
5515
5520
5525
5530
5535
5540
5545
5550
5555
5560
5565
5570
5575
5580
5585
5590
5595
5600
5605
5610
5615
5620
5625
5630
5635
5640
5645
5650
5655
5660
5665
5670
5675
5680
5685
5690
5695
5700
5705
5710
5715
5720
5725
5730
5735
5740
5745
5750
5755
5760
5765
5770
5775
5780
5785
5790
5795
5800
5805
5810
5815
5820
5825
5830
5835
5840
5845
5850
5855
5860
5865
5870
5875
5880
5885
5890
5895
5900
5905
5910
5915
5920
5925
5930
5935
5940
5945
5950
5955
5960
5965
5970
5975
5980
5985
5990
5995
6000
6005
6010
6015
6020
6025
6030
6035
6040
6045
6050
6055
6060
6065
6070
6075
6080
6085
6090
6095
6100
6105
6110
6115
6120
6125
6130
6135
6140
6145
6150
6155
6160
6165
6170
6175
6180
6185
6190
6195
6200
6205
6210
6215
6220
6225
6230
6235
6240
6245
6250
6255
6260
6265
6270
6275
6280
6285
6290
6295
6300
6305
6310
6315
6320
6325
6330
6335
6340
6345
6350
6355
6360
6365
6370
6375
6380
6385
6390
6395
6400
6405
6410
6415
6420
6425
6430
6435
6440
6445
6450
6455
6460
6465
6470
6475
6480
6485
6490
6495
6500
6505
6510
6515
6520
6525
6530
6535
6540
6545
6550
6555
6560
6565
6570
6575
6580
6585
6590
6595
6600
6605
6610
6615
6620
6625
6630
6635
6640
6645
6650
6655
6660
6665
6670
6675
6680
6685
6690
6695
6700
6705
6710
6715
6720
6725
6730
6735
6740
6745
6750
6755
6760
6765
6770
6775
6780
6785
6790
6795
6800
6805
6810
6815
6820
6825
6830
6835
6840
6845
6850
6855
6860
6865
6870
6875
6880
6885
6890
6895
6900
6905
6910
6915
6920
6925
6930
6935
6940
6945
6950
6955
6960
6965
6970
6975
6980
6985
6990
6995
7000
7005
7010
7015
7020
7025
7030
7035
7040
7045
7050
7055
7060
7065
7070
7075
7080
7085
7090
7095
7100
7105
7110
7115
7120
7125
7130
7135
7140
7145
7150
7155
7160
7165
7170
7175
7180
7185
7190
7195
7200
7205
7210
7215
7220
7225
7230
7235
7240
7245
7250
7255
7260
7265
7270
7275
7280
7285
7290
7295
7300
7305
7310
7315
7320
7325
7330
7335
7340
7345
7350
7355
7360
7365
7370
7375
7380
7385
7390
7395
7400
7405
7410
7415
7420
7425
7430
7435
7440
7445
7450
7455
7460
7465
7470
7475
7480
7485
7490
7495
7500
7505
7510
7515
7520
7525
7530
7535
7540
7545
7550
7555
7560
7565
7570
7575
7580
7585
7590
7595
7600
7605
7610
7615
7620
7625
7630
7635
7640
7645
7650
7655
7660
7665
7670
7675
7680
7685
7690
7695
7700
7705
7710
7715
7720
7725
7730
7735
7740
7745
7750
7755
7760
7765
7770
7775
7780
7785
7790
7795
7800
7805
7810
7815
7820
7825
7830
7835
7840
7845
7850
7855
7860
7865
7870
7875
7880
7885
7890
7895
7900
7905
7910
7915
7920
7925
7930
7935
7940
7945
7950
7955
7960
7965
7970
7975
7980
7985
7990
7995
8000
8005
8010
8015
8020
8025
8030
8035
8040
8045
8050
8055
8060
8065
8070
8075
8080
8085
8090
8095
8100
8105
8110
8115
8120
8125
8130
8135
8140
8145
8150
8155
8160
8165
8170
8175
8180
8185
8190
8195
8200
8205
8210
8215
8220
8225
8230
8235
8240
8245
8250
8255
8260
8265
8270
8275
8280
8285
8290
8295
8300
8305
8310
8315
8320
8325
8330
8335
8340
8345
8350
8355
8360
8365
8370
8375
8380
8385
8390
8395
8400
8405
8410
8415
8420
8425
8430
8435
8440
8445
8450
8455
8460
8465
8470
8475
8480
8485
8490
8495
8500
8505
8510
8515
8520
8525
8530
8535
8540
8545
8550
8555
8560
8565
8570
8575
8580
8585
8590
8595
8600
8605
8610
8615
8620
8625
8630
8635
8640
8645
8650
8655
8660
8665
8670
8675
8680
8685
8690
8695
8700
8705
8710
8715
8720
8725
8730
8735
8740
8745
8750
8755
8760
8765
8770
8775
8780
8785
8790
8795
8800
8805
8810
8815
8820
8825
8830
8835
8840
8845
8850
8855
8860
8865
8870
8875
8880
8885
8890
8895
8900
8905
8910
8915
8920
8925
8930
8935
8940
8945
8950
8955
8960
8965
8970
8975
8980
8985
8990
8995
9000
9005
9010
9015
9020
9025
9030
9035
9040
9045
9050
9055
9060
9065
9070
9075
9080
9085
9090
9095
9100
9105
9110
9115
9120
9125
9130
9135
9140
9145
9150
9155
9160
9165
9170
9175
9180
9185
9190
9195
9200
9205
9210
9215
9220
9225
9230
9235
9240
9245
9250
9255
9260
9265
9270
9275
9280
9285
9290
9295
9300
9305
9310
9315
9320
9325
9330
9335
9340
9345
9350
9355
9360
9365
9370
9375
9380
9385
9390
9395
9400
9405
9410
9415
9420
9425
9430
9435
9440
9445
9450
9455
9460
9465
9470
9475
9480
9485
9490
9495
9500
9505
9510
9515
9520
9525
9530
9535
9540
9545
9550
9555
9560
9565
9570
9575
9580
9585
9590
9595
9600
9605
9610
9615
9620
9625
9630
9635
9640
9645
9650
9655
9660
9665
9670
9675
9680
9685
9690
9695
9700
9705
9710
9715
9720
9725
9730
9735
9740
9745
9750
9755
9760
9765
9770
9775
9780
9785
9790
9795
9800
9805
9810
9815
9820
9825
9830
9835
9840
9845
9850
9855
9860
9865
9870
9875
9880
9885
9890
9895
9900
9905
9910
9915
9920
9925
9930
9935
9940
9945
9950
9955
9960
9965
9970
9975
9980
9985
9990
9995
10000
10005
10010
10015
10020
10025
10030
10035
10040
10045
10050
10055
10060
10065
10070
10075
10080
10085
10090
10095
10100
10105
10110
10115
10120
10125
10130
10135
10140
10145
10150
10155
10160
10165
10170
10175
10180
10185
10190
10195
10200
10205
10210
10215
10220
10225
10230
10235
10240
10245
10250
10255
10260
10265
10270
10275
10280
10285
10

Additional selectable marker genes of bacterial origin that confer resistance to antibiotics include gentamycin acetyl transferase, streptomycin phosphotransferase, aminoglycoside-3'-adenyl transferase, the bleomycin resistance determinant. Hayford et al., *Plant Physiol.* 86: 1216 (1988), Jones et al., *Mol. Gen. Genet.*, 210: 86 (1987), Svab et al., *Plant Mol. Biol.* 14: 197 (1990), Hille et al., *Plant Mol. Biol.* 7: 171 (1986). Other selectable marker genes confer resistance to herbicides such as glyphosate, glufosinate or broxynil. Comai et al., *Nature* 317: 741-744 (1985), Gordon-Kamm et al., *Plant Cell* 2: 603-618 (1990) and Stalker et al., *Science* 242: 419-423 (1988).

Other selectable marker genes for plant transformation are not of bacterial origin. These genes include, for example, mouse dihydrofolate reductase, plant 5-enolpyruvylshikimate-3-phosphate synthase and plant acetolactate synthase. Eichholtz et al., *Somatic Cell Mol. Genet.* 13: 67 (1987), Shah et al., *Science* 233: 478 (1986), Charest et al., *Plant Cell Rep.* 8: 643 (1990).

Although many of these markers have been used for selecting transformed plant tissue, these selection systems involving toxic chemical agents can have disadvantages or limitations. One disadvantage is that it may be difficult to recover normal, viable transformed plants directly from chemical selection. Everett et al., *Bio/Technology* 5: 1201-1204 (1987). Another disadvantage is that not all selectable marker systems work for all tissues, in all plant species, due in part to differences in sensitivity of a particular tissue or plant species to the selective agent. The success of any given marker for transformation of a given plant species is not easily predicted. Moreover, potential regulatory issues surrounding the use of antibiotic resistance genes and the use of herbicide resistance genes for plant species capable of outcrossing with weedy species are additional disadvantages of these markers.

Another class of marker genes for plant transformation require screening of presumptively transformed plant cells rather than direct genetic selection of transformed cells for resistance to a toxic substance such as an antibiotic. These genes are particularly useful to quantify or visualize the spatial pattern of expression of a gene in specific tissues and are frequently referred to as reporter genes because they can be fused to a gene or gene regulatory sequence for the investigation of gene expression. Commonly used genes for screening presumptively transformed cells include β -glucuronidase (GUS), β -galactosidase, luciferase, and chloramphenicol acetyltransferase. Jefferson, R.A., *Plant Mol. Biol. Rep.* 5: 387 (1987).., Teeri et al., *EMBO J.* 8: 343 (1989), Koncz et al., *Proc. Natl. Acad. Sci. U.S.A.* 84: 131 (1987), De Block et al., *EMBO J.* 3: 1681 (1984). Another approach to the identification of relatively rare transformation events has been use of a gene that encodes a dominant constitutive regulator of the *Zea mays* anthocyanin pigmentation pathway. Ludwig et al., *Science* 247: 449 (1990).

Although chemical selection of plant cells transformed with selectable marker genes has been successful with plant species and varieties that are easily cultured in vitro, the choice of selectable marker systems that have been shown to be successful for cereals and many other agronomically important plant species is very limited. In general, plant species that tend toward organogenesis and/or shoot propagation have been difficult to transform by means of chemical selection. The success rate with these plant species continues to improve, however, as evidenced by recent advances in Type I selection of maize inbreds and small grain cereals such as barley. Koziel et al., *Bio/Technology* 11: 194-200 (1993) and Mendel et al. In: *Transformation of Plants and Soil Microorganisms*, Wang et al. eds., Cambridge Press (1995).

Likewise, there has been little success in using visual screening methods for primary identification of transformed cells. The GUS gene was used to investigate germline transmission. McCabe et al., *Plant Physiol.*, 87(3): 671 (1988) and McCabe et al., *Plant Cell Tissue Organ Cult.* 33(3): 227 (1993). Histochemical staining for GUS activity was used to locate transgenic sectors in cotton and soybean transformants that ultimately produced transgenic seeds. Since histochemical analysis for GUS activity requires destruction of portions of the presumptively transformed plant tissue, this method is labor intensive and impractical for routine production of transgenic plants. This method is particularly unsuitable for plant species such as maize and other cereals in which transformants are recovered, even under optimum conditions, at low frequency. Recovery of transformed progeny was reported once in barley using GUS expression as a screening tool, but the method was found to be very labor-intensive. Ritala et al., *Plant Mol. Biol.* 24: 317-325 (1994). There have been no reports at all of success with GUS or other screenable markers with maize.

More recently, in vivo methods for visualizing GUS activity that do not require destruction of plant tissue have been made available. Molecular Probes Publication 2908, *Imagene Green™*, p. 1-4 (1993) and Naleway et al., *J. Cell Biol.* 115: 151a (1991). However, these in vivo methods for visualizing GUS activity have not proven useful for recovery of transformed cells because of low sensitivity and high fluorescent backgrounds.

Despite the fact that luciferase genes have been available for many years, this strategy for visualizing transformed cells has also not been successfully adapted for routine recovery of plant transformants. The luciferase-based screening methods are limited by the fact that most of these systems require the presence of luciferin, a compatible luciferase and an exogenously supplied cofactor. In the absence of the substrate,

enzyme or cofactor, the system does not bioluminesce. Cells transformed with a luciferase gene must have cell walls and plasma membranes that are permeable, or rendered permeable, to a compatible luciferin in order to detect bioluminescence. For example, tobacco plants regenerated from cells transformed with a firefly luciferase gene and exposed to a liquid medium containing firefly luciferin exhibited bioluminescence primarily along their major veins. Ow et al., *Science* 234: 856 (1986). Accordingly, a screening method has not been successfully developed for routine plant transformation that does not involve chemical selection or assays, often labor-intensive, that require the sacrifice or destruction of tissue samples for analysis.

A gene encoding GFP has been utilized as a marker for gene expression in prokaryotic and eukaryotic cells. Chalfie et al., *Science* 263: 802 (1994). Many cnidarians utilize GFPs as energy-transfer acceptors in bioluminescence. A gene encoding GFP isolated from a cnidarian and expressed in a heterologous prokaryotic or eukaryotic host produces a protein capable of fluorescence. A cDNA encoding the *Aequorea victoria* GFP produced a fluorescent product when expressed in *Escherichia coli* or *Caenorhabditis elegans* cells. Green fluorescence was detected in transformed cells upon illumination with a long-wave ultraviolet (UV) source without having to supply substrates or cofactors. In addition, fluorescence was stable for at least 10 min when illuminated with 450 to 490 nm light. Transformation of plant cells with a gene encoding GFP and detection of fluorescence has been reported. Haseloff et al., *TIG* 11: 328-329 (1995). Transformed *Arabidopsis* cells could be regenerated into whole plants. However, the regenerated plants expressing GFP exhibited signs of mild to moderate toxicity in the light compared to plants not expressing GFP. The strongest GFP expressors proved more difficult to regenerate. Likewise, in a recent report describing GFP expression in

kanamycin-selected tobacco transformants, Chiu et al.,
Current Biol. 6: 325-330 (1996) note that high GFP
expression levels inhibited regeneration of transgenic
plants.

5 A need therefore exists for a cell transformation
method which does not rely, or does not rely solely on,
selection of cells carrying a gene that confers
resistance to a toxic substance. A need exists for a
method for efficiently and easily identifying transformed
10 plant cells with a visual screenable marker. A need also
exists for a method of cell transformation that does not
require destruction of presumptively transformed tissue
to assay for the presence of a selectable marker gene.
A need exists for a method for cell transformation that
15 combines a selectable marker gene and a screenable marker
gene. In addition, a need exists for a method of cell
transformation which does not require exogenous supply of
a substrate or cofactor for detection of the polypeptide
encoded by a selectable marker gene. Yet another need
20 exists for a method of cell transformation that
circumvents the cellular toxicity of the GFP.

SUMMARY OF THE INVENTION

25 Accordingly, it is an object of the present
invention to provide a method for cell transformation
which does not depend on selection of cells carrying a
gene that confers resistance to a toxic substance.

30 It is another object of the present invention to
provide a method for cell transformation which combines
the selection of cells carrying a gene that confers
resistance to a toxic substance with screening cells for
the presence of a substance that renders transformed
cells identifiable.

35 It is a further object of the present invention to
provide a method for cell transformation which does not
require destruction of presumptively transformed tissue
to assay for the presence of a selectable marker gene.

Yet another object of the present invention is to provide a method for cell transformation which does not require that an exogenous substrate or cofactor be provided to assay for a polypeptide encoded by a selectable marker gene.

It is another object of the present invention to provide a method for cell transformation that circumvents the cellular toxicity of the GFP.

These and other objects are achieved, in accordance with one embodiment of the present invention, by providing an isolated DNA molecule comprising a nucleotide sequence encoding the GFP operably linked to an inducible promoter. The inducible promoter can be selected from the group consisting of the estrogen-inducible promoter, the estradiol-inducible promoter, the ACE1 promoter, the IN2 promoter and the tetracycline repressor promoter.

Also provided is an isolated DNA molecule comprising as nucleotide sequence encoding the GFP wherein the nucleotide sequence is operably linked to a targeting sequence for subcellular localization which directs a protein to a subcellular compartment.

An isolated DNA molecule is provided comprising a nucleotide sequence selected from the group consisting of (a) SEQ ID NO: 1; (b) a nucleotide sequence that has substantial sequence similarity with SEQ ID NO: 1; and (c) a functional fragment of (a) or (b), wherein said DNA molecule encodes a GFP. The nucleotide sequence encoding a GFP may be operably linked to a nucleotide sequence encoding a targeting sequence for subcellular localization which directs a protein to a subcellular compartment. The nucleotide sequence encoding a GFP and a targeting sequence may further comprise a promoter operably linked to said nucleotide sequence.

Also provided are expression vectors comprising DNA molecules encoding a GFP, optionally operably linked to an inducible promoter and/or targeting sequence. The expression vector of the instant invention may carry a

nucleotide sequence encoding a foreign protein operably linked to a second promoter.

Also provided is a method of using the expression vectors of the instant invention to produce a transformed plant, comprising the steps of introducing an expression carrying a gene encoding GFP into regenerable plant cells and selecting cells containing the GFP for regeneration. The method of the instant invention may include the step of inducing GFP expression where the nucleotide sequence encoding GFP is operably linked to an inducible promoter. The regenerable plant cells utilized in the method of the instant invention are selected from the group consisting of *Zea*, *Brassica* and *Helianthus* cells.

Also provided are transgenic plants expressing the isolated DNA molecule encoding GFP. In addition, transgenic plants comprising a vector carrying a nucleotide sequence encoding GFP are provided.

A method for producing a transgenic plant is provided comprising the steps of (a) constructing an expression vector comprising (i) a first promoter which is an inducible promoter, operably linked to a nucleotide sequence encoding a GFP, and (ii) a second promoter operably linked to a foreign gene; (b) introducing said expression vector into regenerable plant cells; (c) inducing expression of the gene encoding the GFP and selecting transformed plant cells containing said protein; and (d) regenerating transformed plants from said selected transformed plant cells. The inducible promoter may be selected from the group including the estrogen-inducible promoter, the estradiol-inducible promoter, the ACE1 promoter, the IN2 promoter and the tetracycline repressor promoter.

Also provided is a method for producing a transgenic plant, comprising the steps: (a) constructing an expression vector comprising (i) a first promoter operably linked to a nucleotide sequence encoding a sequence for subcellular localization which directs a protein to a subcellular compartment which is operably

linked to a nucleotide sequence encoding a GFP, and (ii) a second promoter operably linked to a foreign gene; (b) introducing said expression vector into regenerable plant cells; (c) selecting transformed plant cells containing said GFP; and (d) regenerating transformed plants from said selected transformed plant cells. The targeting sequence for subcellular localization directs the GFP to the mitochondria, chloroplasts, peroxisomes, vacuole, endoplasmic reticulum, cell wall or for secretion generally into the apoplast. This method may include a nucleotide sequence encoding a GFP selected from the group consisting of: (a) SEQ ID NO: 1; (b) a nucleotide sequence that has substantial sequence similarity with SEQ ID NO: 1; and (c) a functional fragment of (a) or (b).

Also provided is a method for producing a transgenic plant, comprising the steps: (a) constructing an expression vector comprising (i) a first promoter operably linked to a nucleotide sequence encoding a screenable marker flanked on the 5-prime and 3-prime ends with a recombinase-specific target sequence, and (ii) a second promoter operably linked to a foreign gene; (b) introducing said expression vector into regenerable plant cells; (c) selecting transformed plant cells containing said screenable marker; (d) transforming the plant cells with a second expression vector containing a gene encoding a site-specific recombinase; (e) selecting plant cells that no longer express the screenable marker; (f) regenerating transformed plants from said selected transformed plant cells; and (g) isolating said foreign protein. The site-specific recombinase may be selected from the group consisting of the FLP/FRT, Ac/DS and cre/lox systems. In addition, the screenable marker may be the GFP.

An isolated DNA molecule comprising a nucleotide sequence encoding the GFP fused in frame to a nucleotide sequence encoding superoxide dismutase is also provided. Cells transformed with the DNA molecule contain a fusion

protein that (i) produces green fluorescence in the presence of UV to blue light and (ii) displays superoxide dismutase activity.

Also provided is a method for producing a transgenic plant comprising: (a) constructing an expression vector comprising (i) a first promoter operably linked to a nucleotide sequence encoding a GFP, and (ii) a second promoter operably linked to a gene encoding an enzyme that is an oxygen scavenger, and (iii) a third promoter operably linked to a foreign gene; (b) introducing said expression vector into regenerable plant cells; (c) selecting transformed plant cells containing said GFP; and (d) regenerating transformed plants from said selected transformed plant cells. The oxygen scavenger enzyme may be superoxide dismutase.

Alternatively, a method for producing a transgenic plant is provided comprising: (a) constructing an expression vector comprising (i) a first promoter operably linked to a nucleotide sequence encoding a fusion protein comprising the GFP and an oxygen scavenger enzyme fused in frame, and (ii) a second promoter operably linked to a foreign gene; (b) introducing said expression vector into regenerable plant cells; (c) selecting transformed plant cells containing said the GFP and oxygen scavenger enzyme activity; and (d) regenerating transformed plants from said selected transformed plant cells. The oxygen scavenger enzyme may be superoxide dismutase.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 presents the nucleotide sequence [SEQ ID NO: 1] encoding a GFP with its corresponding amino acid sequence [SEQ ID NO: 2].

Figure 2 shows the Plasmid Map of PHP167.

Figure 3 shows the Plasmid Map of PHP762.

Figure 4 shows the Plasmid Map of PHP7921.

Figure 5 shows the Plasmid Map of PHP8144.

DETAILED DESCRIPTION

1. Definitions

5 In the description that follows, a number of terms are used extensively. The following definitions are provided to facilitate understanding of the invention.

10 A structural gene is a DNA sequence that is transcribed into messenger RNA (mRNA) which is then translated into a sequence of amino acids characteristic of a specific polypeptide.

15 A promoter is a DNA sequence that directs the transcription of a structural gene. Typically, a promoter is located in the 5-prime region of a gene, proximal to the transcriptional start site of a structural gene. If a promoter is an inducible promoter, then the rate of transcription increases in response to an inducing agent. In contrast, the rate of transcription is not regulated by an inducing agent if the promoter is a constitutive promoter. For example, a promoter may be regulated in a tissue-specific or tissue-preferred manner such that it is only active in transcribing the associated coding region in a specific tissue type(s) such as leaves, roots or meristem.

25 A cytoplasmically localized protein is a protein coded for by a gene which does not include any specific signal for targeting of the protein into any subcellular organelle or compartment or any signals for secretion of the protein. For example, the GFP structural gene operably linked 5-prime to a promoter and to an appropriate 3-prime sequence encodes for a protein compartmentalized in the cytoplasm.

30 A signal or targeting sequence is a structural peptide domain required for targeting of a given polypeptide to a subcellular organelle, subcellular compartment or secretion from the cell. As used herein, the phrase sequence for subcellular localization is

intended to refer collectively to any form of signal, targeting or retention sequence as defined herein. Signal sequences for a polypeptide directed to the chloroplast or mitochondrion are largely localized to the amino-terminus of the polypeptide. Signal sequences for a polypeptide directed to the glyoxysomes and peroxisomes are largely localized to the carboxy-terminal domains of the polypeptide. Accordingly, targeted transport of the GFP protein is accomplished by means of chemically joining in proper reading frame, or operably linking, the nucleotide sequence encoding a signal sequence to the 5-prime and/or 3-prime region of the GFP structural gene.

A mitochondrial targeting sequence facilitates the transport of a protein to a mitochondrial compartment. Typically, the mitochondrial targeting sequence is located at the amino-terminus of a polypeptide.

A secretion targeting sequence targets a polypeptide for export into the extracellular space through the ER. For example, operably linking a nucleotide sequence encoding the barley alpha amylase 1 (BAA) secretory targeting sequence to the 5-prime end of a structural gene targets the encoded protein for export into the extracellular space.

A cell wall targeting sequence targets a polypeptide for export from the cell but the polypeptide is specifically localized to the cell wall. For example, cell wall localization of a polypeptide is accomplished by operably linking a nucleotide sequence encoding BAA 5-prime, and operably linking a nucleotide sequence encoding a portion of the maize hydroxyproline-rich glycoprotein 3-prime to a gene encoding the polypeptide. Steifel et al., Plant Cell 2: 785-793 (1990).

A vacuolar signal sequence facilitates the transport of a protein to the vacuole. For example, vacuolar targeting is accomplished by fusing the BAA secretory signal sequence at the amino-terminus of the protein and a sequence encoding a vacuolar signal sequence to the carboxy-terminus. Transport of a polypeptide to the

vacuole is therefore accomplished by means operably linking a nucleotide sequence encoding BAA 5-prime, and a nucleotide sequence encoding a vacuolar signal sequence 3-prime to a gene encoding a polypeptide. Alternatively, vacuolar targeting is accomplished by constructing a nucleotide sequence comprising in the 5-prime to 3-prime direction nucleotide sequences encoding a vacuole signal sequence, BAA and a polypeptide.

An endoplasmic reticulum retention sequence targets a polypeptide for localization in the lumen of the endoplasmic reticulum. For example, a polypeptide is targeted for retention in the endoplasmic reticulum through the addition of the BAA sequence on the amino-terminus and an endoplasmic reticulum signal sequence on the carboxy-terminus of the polypeptide.

A nuclear targeting sequence facilitates transport of a polypeptide to the nucleus. Typically, the nuclear signal sequence is located at the amino-terminus of a polypeptide. In order to retain the nuclear targeted protein in the nucleus, it may be necessary to increase the molecular weight of the protein by means of fusing an unrelated protein to the carboxy-terminus of the targeted protein. For example, GFP was retained in the nucleus by operably linking a nucleotide sequence encoding a nuclear signal sequence 5-prime and a nucleotide sequence encoding maize acetolactate synthase 3-prime to a gene encoding a polypeptide.

A peroxisomal targeting sequence facilitates the transport of a polypeptide into the peroxisome. Typically, the peroxisomal signal sequence is a tripeptide located at the carboxy-terminus of a polypeptide.

A chloroplast targeting sequence facilitates the transport of a nuclear encoded protein to a chloroplast compartment. Typically, the chloroplast signal sequence is located at the amino-terminus of a polypeptide. Accordingly, transport of a polypeptide to a chloroplast compartment is accomplished by means of operably linking

the nucleotide sequence encoding a chloroplast signal sequence to the 5-prime region of a gene encoding a polypeptide.

5 An isolated DNA molecule is a fragment of DNA that is not integrated in the genomic DNA of an organism.

10 Complementary DNA (cDNA) is a single-stranded DNA molecule that is formed from an mRNA template by the enzyme reverse transcriptase. Typically, a primer complementary to portions of mRNA is employed for the initiation of reverse transcription. Those skilled in the art also use the term "cDNA" to refer to a double-stranded DNA molecule consisting of such a single-stranded DNA molecule and its complementary DNA strand.

15 The term expression refers to the biosynthesis of a gene product. For example, in the case of a structural gene, expression involves transcription of the structural gene into mRNA and the translation of mRNA into one or more polypeptides.

20 A cloning vector is a DNA molecule, such as a plasmid, cosmid, or bacteriophage, that has the capability of replicating autonomously in a host cell. Cloning vectors typically contain one or a small number of restriction endonuclease recognition sites at which foreign DNA sequences can be inserted in a determinable fashion without loss of an essential biological function of the vector, as well as a marker gene that is suitable for use in the identification and selection of cells transformed with the cloning vector. Marker genes typically include, for example, genes that provide tetracycline resistance, ampicillin resistance or
30 kanamycin resistance.

35 An expression vector is a DNA molecule comprising a gene that is expressed in a host cell. Typically, gene expression is placed under the control of certain regulatory elements, including constitutive or inducible promoters, tissue-specific regulatory elements, and enhancers. Such a gene is said to be operably linked to the regulatory elements.

Transformation includes introduction of genetic material into plant cells resulting in chromosomal integration and stable heritability through meiosis. Transformation also includes introduction of genetic material into plant cells in the form of plant viral vectors involving epichromosomal replication and gene expression which may exhibit variable properties with respect to meiotic stability.

A foreign gene refers in the present description to a DNA sequence that is operably linked to at least one heterologous regulatory element.

A recombinant host may be any prokaryotic or eukaryotic cell that contains either a cloning vector or expression vector. This term also includes those prokaryotic or eukaryotic cells that have been genetically engineered to contain the cloned gene(s) in the chromosome or genome of the host cell.

A transgenic plant is a plant having one or more plant cells that contain an expression vector.

A green fluorescent protein (GFP), or a functional fragment of a GFP, is capable of producing a green fluorescence. GFP absorbs in the UV to blue range with a peak at 395 nm and emits in the green with a peak at 510 nm. The "red-shifted" version of GFP is a modified version of GFP which absorbs between 480 to 490 nm and emits in the green with a peak at 510nm. Red-shifted GFP is abbreviated as GFP_r. The "blue fluorescent protein" is a modified version of GFP which absorbs around 380nm and emits in the blue with a peak at around 445nm. Blue fluorescent protein is abbreviated as BFP. GFP_n is a nucleotide sequence coding for GFP in which the DNA sequence has been modified based on codon preference in maize (Fig. 1).

Two nucleic acid molecules are considered to have a substantial sequence similarity if their nucleotide sequences share a similarity of at least 50%. Sequence similarity determinations can be performed, for example, using the FASTA program (Genetics Computer Group;

Madison, WI). Alternatively, sequence similarity determinations can be performed using BLAST (Basic Local Alignment Search Tool) of the Experimental GENIFO(R) BLAST Network Service. See Altschul et al., *J. Mol. Biol.* 215: 403 (1990). Also, see Pasternak et al., "Sequence Similarity Searches, Multiple Sequence Alignments, and Molecular Tree Building," in *METHODS IN PLANT MOLECULAR BIOLOGY AND BIOTECHNOLOGY*, Glick et al. (eds.), pages 251-267 (CRC Press 1993).

2. Reducing GFP Toxicity in Transformed Plants

Oxidative stress associated with GFP fluorescence has been suggested to cause cellular toxicity. Haseloff et al., *TIG* 11: 328-329 (1995). If GFP is to be useful as a screenable marker for plant transformation, as well as a reporter gene for detection of gene expression, strategies for reducing GFP toxicity must be available. The gene encoding GFP can be operably linked to an inducible promoter so that GFP can be transiently expressed and transformed cells identified. Alternatively, the gene encoding GFP can be operably linked to a signal sequence for targeting to an organelle or subcellular compartment or for secretion to the apoplast where GFP is not toxic to the cell.

Oxidative stress can be ameliorated by transforming cells containing GFP with an enzyme that serves as an oxygen scavenger. Such enzymes are well known in the art. For example, genes encoding the enzyme superoxide dismutase may protect against the primary oxidative stresses associated with GFP fluorescence. Balzan et al., *Proc. Nat Acad. Sci.* 92: 4219-4223 (1995). Other enzymes involved in the oxidative stress response, such as ascorbate peroxidase or glutathione S-transferase, could help mitigate secondary effects in the cell. Conklin, *Plant Physiol.* 109: 203-212 (1995).

More specifically, oxidative stress can be ameliorated by transforming cells with a DNA construct

5 carrying genes encoding GFP and SOD. Alternatively, the gene encoding GFP can be fused in frame to the gene encoding SOD. Cells transformed with this DNA construct produce a fusion protein that cause the cells to fluoresce in the presence of UV-blue light and express SOD activity.

10 The toxicity of GFP in transformed plants can be eliminated by excising the screenable marker gene following detection of transformed cells or sectors. The FLP/FRT system is used in conjunction with GFPm as a visible viable marker for FRT excision. The FLP/FRT system has been demonstrated in maize suspension cells using GUS expression as an indicator of FRT excision. Lysnik et al., NAR 21: 969-975 (1993). For example,
15 plant cells are bombarded with a DNA construct containing the GFP gene flanked by FRT sequences as well as a foreign or agronomic gene of interest. The GFP gene may be operably linked to a constitutive promoter or an inducible promoter. In addition, the GFP gene may be
20 operably linked to a signal sequence. Stable transformants are detected by means of screening for GFP.

Transgenic callus pieces are spread on medium and bombarded a second time with a FLP recombinase construct. Callus is monitored periodically under UV to blue
25 illumination to detect cells that no longer express GFP. Callus pieces that no longer express GFP are regenerated and analyzed for expression of the foreign or agronomic gene. Agronomically useful transgenic plants are thereby produced that do not contain a marker gene.

30 The sensitivity of the screening method can be further increased by placing two markers genes between the FRT sequences. For example, plant cells are bombarded with a DNA construct containing the PAT and GFP genes flanked by FRT sequences and a foreign or agronomic
35 gene. Stable transformants are recovered on bialaphos-containing medium and positive GFP-expression is confirmed. The transformed callus is then bombarded with FLP. The callus is then grown for 2-6 weeks with no

selection until clear GFP-null sectors can be identified. These sectors can be transferred onto bialaphos/chlorophenol red multiwell test plates to confirm bialaphos sensitivity (i.e. within 3-5 days). Callus pieces that no longer express GFP or PAT are regenerated and analyzed for expression of the foreign or agronomic gene. This permits recovery of agronomically useful transformants without any marker genes in the final product.

Likewise, the Ac/Ds system of maize can also be used in transgenic plants to excise the screenable marker gene that is transformed together with a foreign or agronomic gene. Mobilization of Ac and/or Ds has been demonstrated in diverse plants such as tomato, tobacco and Arabidopsis. Yoder et al., In Tomato Technology, Alan R. Liss, Inc. pp 189-198 (1987); Yoder et al., U.S. Patent No. 5,225,341; Baker et al., *EMBO J* 6: 1547-1554 (1987) and Lawson et al. *Mol. Gen. Genet.*, 245: 608-615 (1994). Likewise, the cre/lox recombinase system from bacteriophage P1 could also be used in conjunction with GFP. Excision of transgenes in plants using the cre/lox system was first demonstrated in tobacco. Odell et al., *Mol. Gen. Genet.*, 223: 369-378 (1990) and Dale and Ow, *Proc. Natl. Acad. Sci. USA*, 88: 10558-10562 (1991). Similar to the FLP and Ac systems described above, GFP expression provides an efficient, easily scorable phenotype for monitoring excision.

3. Isolation of Nucleotide Sequences Encoding GFP

Peptide sequences from purified GFPs are used to design degenerate oligonucleotide primers for polymerase chain reactions and gene cloning. Protein extracts can be prepared from cnidarian cells by standard methods known to the art. See, for example, Harlow and Lane, *Antibodies, A Laboratory Manual*, Cold Spring Harbor Press, (1988). In a preferred embodiment, cnidarian cells are extracted into a buffer, and the extracts

separated into membrane and soluble fractions. Each fraction is tested for green fluorescence activity in the presence of blue light.

5 Fractions which contain GFP activity are then purified further to determine which components are responsible for the activity. Purification of the active fractions can be carried out by methods known in the art. See, for example, PROTEIN PURIFICATION METHODS - A PRACTICAL APPROACH, Harris et al., Eds. (IRL Press, 10 Oxford, 1989). Extracts prepared as described above are purified by sequential size exclusion chromatography, isoelectric focusing, HPLC size exclusion chromatography, and chromatography on an affinity column. Fractions which display GFP activity can be analyzed further by 15 SDS-PAGE analysis to determine the approximate molecular mass of the active component.

Purified GFPs prepared by the methods described above can be sequenced using methods well known in the art, for example using a gas phase peptide sequencer 20 (Applied Biosystems, Foster City, CA). To determine as much of the peptide sequence as possible, it is preferred that the proteins of the present invention be cleaved into smaller fragments more suitable for gas-phase sequence analysis. This can be achieved by treatment of 25 a purified GFP with selective peptidases, and in a particularly preferred embodiment, with endoproteinase lys-C (Boehringer). The fragments so produced can be separated by reversed-phase HPLC chromatography.

30 The peptide sequences of the proteins determined as above can be used to determine the DNA sequence encoding the protein. Methods for carrying out this determination are well known in the art. See, for example Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, Second Edition, (Cold Spring Harbor Laboratory Press, Cold 35 Spring Harbor, NY 1989).

In a preferred embodiment of the present invention, the peptide sequences are used to design degenerate oligonucleotide primers for polymerase chain reactions.

Each degenerate primer set will preferably contain every possible DNA sequence encoding the corresponding peptide sequences. Primer sets are prepared in both the sense and antisense orientation. Suitable oligonucleotide primers can be synthesized using commercial synthesizers, such as those supplied by Applied Biosystems (Foster City, CA). In a particularly preferred embodiment, the primers include additional nucleotide sequences containing restriction endonuclease cleavage sites. The presence of such sites allows for the directional cloning of PCR products into suitable cloning vectors after treatment with an appropriate restriction enzyme. See Finney, "Molecular Cloning of PCR Products" in CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, Ausubel et al. Eds. (John Wiley & Sons, New York, 1987) p. 15.7.1.

Template DNA for the PCR can be prepared from appropriate cnidarian cells or tissues using methods well known in the art. See Sambrook et al., *supra*. In a preferred embodiment, host cells are crushed under liquid nitrogen and mRNA is extracted using a commercially available kit (Pharmacia, Piscataway, NJ).

The mRNA preparation can then be used as a template for cDNA synthesis using poly(dT) or random hexamer primers by standard techniques. See Sambrook et al., *supra*. In a particularly preferred embodiment, cDNA synthesis is carried out using a commercially available kit (Pharmacia).

The cDNA can then be used directly for PCR using the method of Saiki et al., *Science* 239: 487 (1988). The cDNA also is used to prepare a cDNA library by standard methods. See Sambrook et al., *supra*. In a particularly preferred embodiment, the cDNA is packaged into bacteriophage particles using a commercially available kit (Promega, Madison, WI). The packaged cDNA is then transfected into *E. coli* to produce a cDNA library.

In an alternative preferred embodiment, genomic DNA from cnidarian cells or tissue can be used as the template DNA for the PCR. Genomic DNA can be prepared by

standard methods and PCR can then be used to prepare double stranded DNA molecules to probe the cDNA library and the genomic DNA for the gene(s) encoding GFP. In a preferred embodiment, degenerate primers are prepared corresponding to the termini of the longest peptide sequence determined by peptide sequencing. In a particularly preferred embodiment, primers are used in a PCR with first strand cDNA as template to amplify the DNA encoding the peptide. PCR is carried out under standard conditions. See Sakai et al., *supra*.

PCR amplification products are analyzed by polyacrylamide gel electrophoresis using standard methods. If an amplification product of the expected size (based on the peptide sequence) is found, the product is digested with appropriate restriction enzymes, ligated into a cloning vector and cloned by standard methods. See Sambrook et al, *supra*. In a preferred embodiment, clones are sequenced to verify that sequences according to the expected peptide sequence are present.

Once the DNA sequence encoding the peptide is known, it can be used to prepare non-degenerate primers corresponding to that sequence, again containing restriction enzyme recognition sequences to aid in cloning of DNA products. These primers are used in combination with degenerate primers corresponding to other peptide sequences to generate PCR amplification products which can be cloned and then analyzed as above. By these means, fragments of the gene sequence of the protein can be determined. Alternative methods for carrying out this PCR analysis include use of the 5' or 3' RACE methods using commercially available kits, such as those manufactured by Life Technologies (Gaithersburg, MD) or Clontech (Palo Alto, CA). Primers for this method are selected according to the manufacturer's directions.

Gene fragments prepared as above are excised from the cloning vector by restriction enzyme digestion, labeled with ³²P by conventional methods and used as probes to identify the complete gene encoding the

protein from within a cDNA or genomic library. In a preferred embodiment, the probe is chosen such that it is long enough to ensure hybridization specificity, while remaining short enough to allow reasonable rates of hybridization to the target gene.

A cnidarian genomic DNA library can be prepared by means well-known in the art. See, for example, Slightom et al. "Construction of λ Clone Banks," in METHODS IN PLANT MOLECULAR BIOLOGY AND BIOTECHNOLOGY, Glick et al. (eds.), pages 121-146 (CRC Press, 1993). Genomic DNA can be isolated from cnidarian tissue, for example, by lysing plant tissue with the detergent Sarkosyl, digesting the lysate with proteinase K, clearing insoluble debris from the lysate by centrifugation, precipitating nucleic acid from the lysate using isopropanol, and purifying resuspended DNA on a cesium chloride density gradient. See, for example, Ausubel et al. (eds.), CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, pages 2.3.1-2.3.3 (Wiley Interscience 1990) ["Ausubel"].

DNA fragments that are suitable for the production of a genomic library can be obtained by the random shearing of genomic DNA or by the partial digestion of genomic DNA with restriction endonucleases. See, for example, Ausubel at pages 5.3.2-5.4.4, and Slightom et al., supra. Genomic DNA fragments can be inserted into a vector, such as a bacteriophage or cosmid vector, in accordance with conventional techniques, such as the use of restriction enzyme digestion to provide appropriate termini, the use of alkaline phosphatase treatment to avoid undesirable joining of DNA molecules, and ligation with appropriate ligases. Techniques for such manipulation are disclosed by Slightom et al., supra, and are well-known in the art. Also see Ausubel at pages 3.0.5-3.17.5.

Screening of the cDNA or genomic library is carried out by conventional methods. See Sambrook et al, supra. Clones which hybridize to the probe are purified and their sequences determined. To facilitate sequencing,

5 nested deletions in the clones can be created using standard protocols, or by commercially available kits such as Erase-a-base (Promega, Madison, WI) or The Deletion Factory (Life Technologies, Gaithersburg, MD), following the manufacturer's instructions. The sequences obtained are analyzed for the presence of open reading frames by conventional methods and to check if the entire gene sequence has been found. In a preferred embodiment, cDNA libraries are prepared by both random hexamer and poly (dT) priming from samples, and are used to maximize the chances of finding the complete coding sequence of the desired gene.

10 Once the entire coding sequence of the gene for GFP has been determined, the gene can be inserted into an appropriate expression system. The gene can be expressed in any number of different recombinant DNA expression systems to generate large amounts of protein, which can then be purified.

20 4. Chemical Synthesis of Genes Encoding GFP

Once the amino acid sequence of a GFP is known, a gene encoding this GFP can be synthesized. In addition, the nucleotide sequence of a synthetic gene encoding GFP can be optimized for expression in plants by modifying the codon usage to include plant preferred codons. See, for example, Murray et al., *NAR* 17: 477 (1989). Even more specifically, the nucleotide sequence of a synthetic gene encoding GFP can be optimized for expression in monocotyledonous or dicotyledonous plants. See, for example, Campbell et al., *Plant Physiol.* 92: 1 (1990).

30 Genes encoding GFP can be obtained, for example, by synthesizing the genes with mutually priming long oligonucleotides. See, for example, Ausubel at pages 8.2.8 to 8.2.13. Also, see Wosnick et al., *Gene* 60:115 (1987). Moreover, current techniques using the polymerase chain reaction provide the ability to synthesize genes as large as 1.8 kilobases in length.

Adang et al., *Plant Molec. Biol.* 21: 1131 (1993); Bambot et al., *PCR Methods and Applications* 2: 266 (1993).

5 5. Identification of Functional Fragments of the GFP

 DNA clones can be analyzed using a variety of techniques such as restriction analysis, Southern analysis, primer extension analysis, and DNA sequence analysis. Primer extension analysis or S1 nuclease protection analysis, for example, can be used to localize the putative start site of transcription of the cloned gene. Ausubel at pages 4.8.1-4.8.5; Walmsley et al., "Quantitative and Qualitative Analysis of Exogenous Gene Expression by the S1 Nuclease Protection Assay," in METHODS IN MOLECULAR BIOLOGY, VOL. 7: GENE TRANSFER AND EXPRESSION PROTOCOLS, Murray (ed.), pages 271-281 (Humana Press Inc. 1991). Functional fragments of the GFP protein are identified by production of green fluorescence in the presence of blue UV light.

 The general approach of such functional analysis involves subcloning DNA fragments of a genomic clone, cDNA clone or synthetic gene sequence into an expression vector, introducing the expression vector into a heterologous host, and screening to detect green fluorescence in the presence of UV to blue light. Methods for generating fragments of a cDNA or genomic clone are well-known. Variants of an isolated DNA encoding GFP can be produced by deleting, adding and/or substituting nucleotides for the isolated nucleotides, for example, the nucleotide sequence of SEQ ID NO: 1. Such variants can be obtained, for example, by oligonucleotide-directed mutagenesis, linker-scanning mutagenesis, mutagenesis using the polymerase chain reaction, and the like. See, for example, Ausubel, pages 8.0.3-8.5.9. Also see generally, McPherson (ed.), DIRECTED MUTAGENESIS: A PRACTICAL APPROACH, (IRL Press 1991). Thus, the present invention also encompasses DNA molecules comprising nucleotide sequences that have

substantial sequence similarity with SEQ ID NO: 1 and encode a GFP.

6. Methods for Plant Transformation

5 The GFP-based screening method for plant transformation of the instant application can be used in conjunction with any method of plant transformation and regeneration. Numerous methods for plant transformation
10 have been developed, including biological and physical, plant transformation protocols. See, for example, Miki et al., "Procedures for Introducing Foreign DNA into Plants" in *Methods in Plant Molecular Biology and Biotechnology*, Glick, B.R. and Thompson, J.E. Eds. (CRC Press, Inc., Boca Raton, 1993) pages 67-88. In addition,
15 expression vectors and in vitro culture methods for plant cell or tissue transformation and regeneration of plants are available. See, for example, Gruber et al., "Vectors for Plant Transformation" in *Methods in Plant Molecular Biology and Biotechnology*, Glick, B.R. and Thompson, J.E. Eds. (CRC Press, Inc., Boca Raton, 1993) pages 89-119.
20

A. Agrobacterium-mediated Transformation

25 The most widely utilized method for introducing an expression vector into plants is based on the natural transformation system of *Agrobacterium*. See, for example, Horsch et al., *Science* 227:1229 (1985). *A. tumefaciens* and *A. rhizogenes* are plant pathogenic soil bacteria which genetically transform plant cells. The Ti and Ri plasmids of *A. tumefaciens* and *A. rhizogenes*,
30 respectively, carry genes responsible for genetic transformation of the plant. See, for example, Kado, C.I., *Crit. Rev. Plant. Sci.* 10: 1 (1991). Descriptions of *Agrobacterium* vector systems and methods for
35 *Agrobacterium*-mediated gene transfer are provided by Gruber et al., *supra*, Miki et al., *supra*, and Moloney et al., *Plant Cell Reports* 8: 238 (1989).

B. Direct Gene Transfer

Despite the fact the host range for *Agrobacterium*-mediated transformation is broad, some major cereal crop species and gymnosperms have generally been recalcitrant to this mode of gene transfer, even though some success has recently been achieved in rice. Hiei et al., *The Plant Journal* 6: 271-282 (1994). Several methods of plant transformation, collectively referred to as direct gene transfer, have been developed as an alternative to *Agrobacterium*-mediated transformation.

A generally applicable method of plant transformation is microprojectile-mediated transformation wherein DNA is carried on the surface of microprojectiles measuring 1 to 4 μm . The expression vector is introduced into plant tissues with a biolistic device that accelerates the microprojectiles to speeds of 300 to 600 m/s which is sufficient to penetrate plant cell walls and membranes. Sanford et al., *Part. Sci. Technol.* 5: 27 (1987), Sanford, J.C., *Trends Biotech.* 6: 299 (1988) Sanford, J.C., *Physiol. Plant* 79: 206 (1990), Klein et al., *Biotechnology* 10: 268 (1992).

Another method for physical delivery of DNA to plants is sonication of target cells. Zhang et al., *Bio/Technology* 9: 996 (1991). Alternatively, liposome or spheroplast fusion have been used to introduce expression vectors into plants. Deshayes et al., *EMBO J.*, 4: 2731 (1985), Christou et al., *Proc Natl. Acad. Sci. U.S.A.* 84: 3962 (1987). Direct uptake of DNA into protoplasts using CaCl_2 precipitation, polyvinyl alcohol or poly-L-ornithine have also been reported. Hain et al., *Mol. Gen. Genet.* 199: 161 (1985) and Draper et al., *Plant Cell Physiol.* 23: 451 (1982). Electroporation of protoplasts and whole cells and tissues have also been described. Donn et al., In Abstracts of VIIth International Congress on Plant Cell and Tissue Culture IAPTC, A2-38, p 53 (1990); D'Halluin et al., *Plant Cell* 4: 1495-1505 (1992) and Spencer et al., *Plant Mol. Biol.* 24: 51-61 (1994).

5 A preferred method is microprojectile-mediated bombardment of immature embryos. The embryos can be bombarded on the embryo axis side to target the meristem at a very early stage of development or bombarded on the scutellar side to target cells that typically form callus and somatic embryos. Targeting of the scutellum using projectile bombardment is well known to those in the art of cereal tissue culture. Klein et al., *Bio/Technol.*, 6: 559-563 (1988); Kartha et al., *Plant Cell Rep.* 8: 429-432
10 (1989); Sautter et al., *Bio/Technol.*, 9: 1080-1085 (1991); Chibbar et al., *Genome*, 34: 435-460 (1991). The scutellar origin of regenerable callus from cereals is well known. Green et al., *Crop Sci.*, 15: 417-421 (1975); Lu et al., *TAG* 62: 109-112 (1982); and Thomas and Scott,
15 *J. Plant Physiol.* 121: 159-169 (1985). Targeting the scutellum and then using chemical selection to recover transgenic plants is well established in cereals. D/Halluin et al., *Plant Cell* 4: 1495-1505 (1992); Perl et al., *MGG* 235: 279-284 (1992); Cristou et al.,
20 *Bio/Technol.* 9: 957-962 (1991). This literature reports DNA targeting of the scutellum and recovery of transgenic callus, plants and progeny based on a chemical selection regime. None of these references teach successful plant transformation wherein transformed cells are: visualized
25 with a screenable marker such as GUS.

A preferred transformation method involves bombardment of the scutellar surface of immature embryos to introduce a fGP expression cassette and any other cotransformed genes. Embryos can be pretreated for 1 to
30 48 hours with a high osmoticum medium or left on a high-osmoticum medium for 24-48 hours after bombardment to improve cell survival and transformation frequencies. Immature embryos are then cultured on typical callus-inducing medium with no selective agent. At each
35 subculture transfer, i.e., every two weeks, the culture is monitored using UV-blue light for GFP fluorescence. Fluorescing calli are separated from non-fluorescing

callus, and grown to the point where plants can be regenerated through standard media progressions.

Many exemplary variations, in terms of target cells and tissue culture routes are known in the art that could be used with GFP in this fashion. Some of these variations include introduction of GFP expression into protoplasts, suspension cells, type II callus, and type I callus, as in Morocz et al., *Theor. Appl. Genet.* 80: 721-726 (1990), Gordon-Kamm et al., *Plant Cell* 2: 603-618 (1990), Fromm et al., *Bio/Technol.* 8: 833-839 (1990) and D'Halluin et al., *Plant Cell* 4: 1495-1505 (1992). Microspores and microspore-derived embryogenic callus also represent a feasible alternative target/culturing route for using GFP screening to recover transformants. See Sun et al., *Plant Cell Rep.* 8: 313-316 (1989) and Mitchell et al., *J. Plant Physiol.* 37: 530-536 (1991). Furthermore, it is expected that the method of the present invention is also be applicable to any dicot species that is known to have a callus stage, for example, tobacco, canola or soybean.

In the case of the maize meristem target, the method entails selective enlargement of transgenic sectors, toward genetic homogeneity, in cell layers that contribute to germline transmission. This method is described in copending U.S. patent application 08/483,091, incorporated herein by reference, and comprises the steps of (A) introducing foreign DNA into a meristem that is not enclosed by sheathing leaves; (B) inducing reorganization of the meristem to increase transgenic sector size, whereby the likelihood that a transgenic sector will contribute to germline transmission is increased; and (C) exposing the meristem to conditions under which it differentiates to form a plantlet, wherein the plantlet contains the transgenic sector or is homogeneously transformed by the foreign DNA, such that the plantlet can be grown into a transformed cereal plant that will transmit the foreign DNA to progeny. The foreign DNA can be introduced into

a plurality of meristems, at least some of which differentiate in step (C) to form a plurality of plantlets.

5 In one preferred embodiment, reorganization is effected through at least one manipulation selected from the group consisting of (i) imposition of a nonlethal selective pressure on the meristems, (ii) mechanically-induced meristem reorganization, and (iii) hormonally-induced shoot multiplication. In another preferred
10 embodiment the conditions in step (C) are such that the meristems undergo maturation and plant differentiation to form shoot apices, and the method further comprises effecting reorganization of meristem tissue in the shoot apices to enlarge transformed sectors or to produce
15 periclinal L2 chimeras. The reorganization in this regard can be effected, for example, by exposing the shoot apices to nonlethal selection pressure such that transformed cells have a competitive growth advantage over nontransformed cells in the shoot apices, and the
20 proportion of transformed cells in the shoot apices is increased. In yet another preferred embodiment, the method further comprises a step before step (B), e.g., before step (A), of wounding the apical dome selectively. A method of the present invention also can comprise the
25 further steps of (i) dissecting out an axillary bud from above the base of a leaf of a plantlet when a chimeric sector is observed in a substantial portion of the leaf, and then (ii) germinating the axillary bud into a whole plant or subjecting the axillary bud to shoot
30 multiplication.

Likewise, the introduction of GFP-encoding DNA into
dicot meristems provides an efficient method of mapping
transgenic sectors and following these up into a germline
and into progeny. Using mature imbibed seeds as the
35 starting material, the meristem is exposed by removal of the cotyledons, and DNA is introduced into meristem cells. A preferred method for introducing DNA into meristem cells is *Agrobacterium*-mediated delivery, but

other DNA delivery methods can be used with this explant such as bombardment with DNA-coated particles, or silicon carbide fiber-mediated delivery. Kaeppler et al., *Plant Cell Rep.* 9: 415-418 (1990). After DNA introduction, the meristem explants are cultured.

Plants can be manipulated, for example, by removal of the apical meristem, to stimulate axillary or secondary buds which can exhibit larger transgenic sectors relative to the primary shoot. Flowers above transgenic shoots are pollinated and the progeny are analyzed for transgene presence and expression. A variety of starting explants can regenerate shoots in sunflower, and thus represent alternative targets for GFP-encoding DNA delivery and transmission to progeny. These include the seedling meristem (as above), also the seedling hypocotyl, the mature cotyledon, the immature cotyledon, zygotic immature embryos, somatic embryos, and primary leaflets. See for example, respectively, Greco et al., *Plant Sci. Lett.* 36: 73-77 (1984); Krauter et al., *Helia* 14: 117-122 (1991); Power Am. J. Bot. 74: 497-503 (1987); Krauter et al., *Theor. Appl. Genet.* 82: 521-525 (1991); Finer, *Plant Cell Rep.* 6: 372-374 (1987), and Greco et al., *Plant Sci. Lett.* 36: 73-77 (1984).

In similar fashion, GFP expression could be used in other dicot species, for example, soybean, to map sectors and/or identify transformants and follow the transgenes through to progeny. A GFP-encoding cassette and other cotransformed genes can be introduced into soybean cotyledonary node cells using *Agrobacterium*. The explants are cultured on Gamborg (B5) medium with or without kanamycin selection. With no selection, GFP expression is used to identify transgenic chimeric or homogeneously transformed plantlets, which are rooted, grown to maturity and pollinated in the greenhouse. Progeny seed are collected for analysis. The use of cotyledonary nodes and *Agrobacterium* is presented by way of example, but other target cells and culturing methods could be used with GFP for soybean. In addition to the

cotyledonary node cells, mature cotyledons and immature cotyledons have both been used for transformation. See Hinchee et al., *Bio/Technol.* 6: 915-922 (1988) and Parrott et al., *Plant Cell Rep.* 7: 615-617 (1989).
5 Meristems from immature soybean seed embryonic axes have also been used for bombardment-mediated soybean transformation. See Christou et al., *Plant Physiol.* 87: 671-674 (1988) and McCabe et al., *Bio/Technol.* 6: 923-926 (1988).

10 7. Promoters

A. Inducible Promoters

15 An inducible promoter is operably linked to a nucleotide sequence encoding a GFP. Optionally, the inducible promoter is operably linked to a nucleotide sequence encoding a signal sequence which is operably linked to a nucleotide sequence encoding GFP. With an inducible promoter the rate of transcription increases in
20 response to an inducing agent.

Any inducible promoter can be used in the instant invention. See Ward et al. *Plant Mol. Biol.* 22: 361-366 (1993). Exemplary inducible promoters include that from the ACE1 system which responds to copper (Mett et al. *PNAS* 90: 4567-4571 (1993)); In2 gene from maize which
25 responds to benzenesulfonamide herbicide safeners (Hershey et al., *Mol. Gen. Genetics* 227: 229-237 (1991) and Gatz et al., *Mol. Gen. Genetics* 243: 32-38 (1994)) or Tet repressor from Tn10 (Gatz et al., *Mol. Gen. Genet.* 227: 229-237 (1991)). A particularly preferred inducible
30 promoter is a promoter that responds to an inducing agent to which plants do not normally respond. An exemplary inducible promoter is the inducible promoter from a steroid hormone gene the transcriptional activity of
35 which is induced by a glucocorticosteroid hormone. Schena et al., *Proc. Natl. Acad. Sci. U.S.A.* 88: 10421 (1991).

The expression vector comprises an inducible promoter operably linked to a nucleotide sequence encoding GFP. The expression vector is introduced into plant cells and presumptively transformed cells are exposed to an inducer of the inducible promoter. The cells are screened for the presence of GFP protein by means of illuminating the cells with UV to blue light and screening for the presence of green fluorescence.

B. Tissue-specific or Tissue Preferred Promoters

A tissue-specific promoter is operably linked to a nucleotide sequence encoding a GFP. Optionally, the tissue-specific promoter is operably linked to a nucleotide sequence encoding a signal sequence which is operably linked to a nucleotide sequence encoding GFP. Plants transformed with a gene encoding GFP operably linked to a tissue-specific promoter produce the GFP protein exclusively, or preferentially, in a specific tissue.

Any tissue-specific or tissue-preferred promoter can be utilized in the instant invention. Exemplary tissue-specific or tissue-preferred promoters include a root-preferred promoter such as that from the phaseolin gene (Murai et al., *Science* 23: 476-482 (1983) and Sengupta-Gopalan et al., *Proc. Natl. Acad. Sci. USA* 82: 3320-3324 (1985)); a leaf-specific and light-induced promoter such as that from *cab* or *rubisco* (Simpson et al., *EMBO J.* 4(11): 2723-2729 (1985) and Timko et al., *Nature* 318: 579-582 (1985)); an anther-specific promoter such as that from *LAT52* (Twell et al., *Mol. Gen. Genet.* 217: 240-245 (1989)); a pollen-specific promoter such as that from *Zm13* (Guerrero et al., *Mol. Gen. Genet.* 224: 161-168 (1993)) or a microspore-preferred promoter such as that from *apγ* (Twell et al., *Sex. Plant Reprod.* 6: 217-224 (1993)).

The expression vector comprises a tissue-specific or tissue-preferred promoter operably linked to a nucleotide sequence encoding GFP. The expression vector is

introduced into plant cells. The cells are screened for the presence of GFP protein by means of illuminating the cells with UV to blue light and screening for the presence of green fluorescence.

5

C. Constitutive Promoters

A constitutive promoter is operably linked to a nucleotide sequence encoding a GFP or the constitutive promoter is operably linked to a nucleotide sequence encoding a signal sequence which is operably linked to a nucleotide sequence encoding GFP.

10

15

20

25

30

35

Many different constitutive promoters can be utilized in the instant invention. Exemplary constitutive promoters include the promoters from plant viruses such as the 35S promoter from CaMV (Odell et al., *Nature* 313: 810-812 (1985) and the promoters from such genes as rice actin (McElroy et al., *Plant Cell* 2: 163-171 (1990)); ubiquitin (Christensen et al., *Plant Mol. Biol.* 12: 619-632 (1989) and Christensen et al., *Plant Mol. Biol.* 18: 675-689 (1992)); pEMU (Last et al., *Theor. Appl. Genet.* 81: 581-588 (1991)); MAS (Velten et al., *EMBO J.* 3: 2723-2730 (1984)) and maize H3 histone (Lepetit et al., *Mol. Gen. Genet.* 231: 276-285 (1992) and Atanassova et al., *Plant Journal* 2(3): 291-300 (1992)).

The ALS promoter, a XbaI/NcoI fragment 5-prime to the *Brassica napus* ALS3 structural gene (or a nucleotide sequence that has substantial sequence similarity to said XbaI/NcoI fragment), represents a particularly useful constitutive promoter. Co-pending Pioneer Hi-Bred International US Patent Application 08/409,297.

The expression vector comprises a constitutive promoter operably linked to a nucleotide sequence encoding GFP. The expression vector is introduced into plant cells and presumptively transformed cells are screened for the presence of GFP protein by means of illuminating the cells with UV to blue light and screening for the presence of green fluorescence.

8. Signal Sequences For Targeting Proteins to Subcellular Compartments

Transport of GFP to a subcellular compartment such as the chloroplast, vacuole, peroxisome, glyoxysome, cell wall or mitochondrion, or for secretion into the apoplast, is accomplished by means of operably linking the nucleotide sequence encoding a signal sequence to the 5-prime and/or 3-prime region of a gene encoding GFP. Targeting sequences at the 5-prime and/or 3-prime end of the structural gene may determine, during protein synthesis and processing, where the encoded protein is ultimately compartmentalized. The presence of a signal sequence directs a polypeptide to either an intracellular organelle or subcellular compartment or for secretion to the apoplast.

In addition to the targeting sequences described above, the addition of amino acids to the encoded protein (fusing GFP to either another protein, a protein fragment, or a peptide) can further influence the fate of GFP. For example, a nuclear localization signal (NLS) alone is not sufficient to accumulate GFP in the nucleus. The GFP protein is sufficiently small that it freely diffuses from the nucleus into the cytoplasm, probably through the nuclear pores. The fusion of GFP to a sufficiently large protein permits accumulation in the nucleus if the GFP protein also contains an NLS at the amino-terminus. GFP can also be targeted to the nucleus by fusing GFP to proteins that are normally targeted to the nucleus.

The following exemplary signal sequences were utilized for targeting of GFP. However, the invention encompasses use of any signal sequence, or combination of signal sequences, to facilitate transport of GFP to a subcellular compartment or the apoplast where the protein is not toxic to the cell but can be detected.

More specifically, GFP is expressed in plant cells transformed with a expression vector carrying a

nucleotide sequence encoding a signal sequence(s), which directs a protein to a subcellular compartment or to the apoplast, operably linked to a gene encoding GFP. This GFP gene construct can be expressed from a constitutive, tissue-specific, tissue-preferred or inducible promoter. GFP is transported to a subcellular compartment or the apoplast where it is not toxic to the plant cell. Cells illuminated with UV to blue light can be detected because they emit green fluorescence. The transformed cells are selected for regeneration. If the expression vector also carries a gene encoding a foreign protein or agronomically useful protein, the regenerated plant can be used for production of this protein.

A. PHP8080 - Ubi::CTP-GFPm

A chloroplast signal sequence from the maize cab-m7 has been characterized and the nucleotide sequence is shown below. Becker et al., *Plant Mol. Biol.* 20: 49 (1992). The signal sequence includes 21 amino acids of the cab-m7 protein that extends beyond the processing site (marked by *).

```

ATG GCG TCC TCC ACG ATG GCC CTC TCC TCC ACC GCC TTC GCC
M  A  S  S  T  M  A  L  S  S  T  A  F  A
-
GGC AAG GCC GTG AAC GTG CCG TCG TCG TCC GCC TTC GAG
G  K  A  V  N  V  P  S  S  S  A  F  E

GCC CGC GTG ACC ATG AGG AAG ACG GCG GCG AAG GCC AAG
A  R  V  T  *M  R  K  T  A  A  K  A  K  K

CCA GCT GCG GCG TCC GGG AGC CCG TGG TAC GGC CCC ATG
P  A  A  A  S  G  S  P  W  Y  G  P  M

```

B. PHP8087 - CPN60::GFPm

The mitochondrial signal sequence from the maize chaperonin 60 gene is shown below. Close, P.S., Master's Theses, Iowa State University (1993). The signal sequence includes intron 1 of cpn60II (marked by -) and

includes 2 amino acids from the cpn60II protein beyond the processing site (marked by *).

5
 ATG TAC CGC GCC GCC GCT AGC CTC GCC TCC AAG GCG CGG CAA GCC GGG AGC
 M Y R A A A S L A S K A R Q A G S
 AGC TCC GCC GCT CGC CAG-gtgaagcagactcgtgtttatcacgcgtgtgatgggtctgatg
 S S A A R Q
 gaggaatcccgccctcagatttggtaatgttgccctgggggtcttagtagttctgctgcagggctgtg
 10 ggtttgctgcatgggtgctgtttatgttggtggcgacttgcgggaatctgtagttcgttcgcgcgcaaaa
 tctaaagctagctcgttaattggcgtactggcgtgggggtttcacattaatctacgggtggtgaactcgtc
 actaccgtcctccagtttagctgttagacaccgaatacactgattggcagttgagaaacattgatctg
 atccagcagaaatcgatgtcttctgtaaatctgttattttattgtcgtgtaaccttggggcagcggcag
 15 tctctaattgatccacgcactcacctctgtgtgtgtgatgctttatag-GTT GGA AGC AGG CTT
 V G S R L
 GCC TGG AGC AGG AAC TAT GCT GCC ATG
 A W S R N Y *A A M

20

C. PHP8144 - Ubi::BAA-GFPm - Cloned sequence includes 1 amino acid beyond processing site (marked by *, lower case bases show linker sequence).

25 The barley alpha amylase 1 signal sequence is shown below. Knox, C., et al., "Structure and Organization of Two Divergent Alpha-Amylase Genes From Barley", Plant Mol. Biol. 9:3-17 (1987). The cloned sequence
 30 includes sequence only up to processing site.

ATG GCC AAC AAG CAC CTG AGC CTC TCC CTC TTC CTC GTG CTC CTC GGC CTC TCC
 M A N K H L S L S L F L V L L G L S
 GCC TCC CTC GCC TCC GGA tcc ATG
 35 A S L A S *G S M

D. PHP8757 - Ubi::BAA-GFPm-BLVT - Signal sequence starts at *, lower case bases show linker sequence.

40 The barley lectin vacuole signal sequence is shown below. Lerner et al., Plant Physiol. 91: 124-129 (1989).

45 AAG atc tac GTG TTC GCC GAG GCC ATC GCC GCC AAC TCC ACC CTC GTG GCC GAG
 K I Y *V F A E A I A A N S T L V A E

E. PHP8758 - Ubi::BAA-GFPm-HDEL - Signal sequence starts at *, lower case bases show linker sequence.

The HDEL endoplasmic reticulum retention signal from maize b-70 is shown below. Fontes et al., Plant Cell 3: 483-496 (1991).

AAG atc tac GAC GGC GGC GTG GAC GAC GAC CAC GAC GAG CTC
K I Y *D G G V D D D H D E L

F. PHP8759 - Ubi::BAA-SVT-GFPm - Marked by ^, lower case bases show linker sequence.

The sweet potato sporamin vacuole signal sequence is shown below. Matsuoka et al., Proc. Natl. Acad. Sci. 88: 834 (1991). The sequence shown includes 4 additional sporamin amino acids beyond the processing site (marked by *).

ATG GCC AAC AAG CAC CTG AGC CTC TCC CTC TTC CTC GTG CTC CTC GGC CTC TCC
M A N K H L S L S L F L V L L G L S
GCC TCC CTC GCC TCC GGA CAC AGC AGG TTC AAC CCC ATC AGG CTG CCC ACC ACC
A S L A S G *H S R F N P I R L P T T
CAC GAG CCC GCC AGC AGC GAG ACC gga tcc ATG
H E P A *S S E T G S M

G. PHP8882 - Ubi::GFPm-SKL - (marked by *)

The consensus peroxisome signal sequence designated SLK is shown below. Gould et al., J. Cell Biol 108: 1657 (1989).

AAG TCG AAG CTT
K *S K L

H. Ubi::GR-GFPm

The pea glutathione reductase mitochondrial and chloroplast signal sequence is shown below. Creissen et al., Plant J. 2: 129 (1991). The sequence shown includes 20 additional amino acids beyond the putative processing site (marked by *).

ATG AAC CAG GCG ATG GCC ACC CCG CTG TCC CTG TCC TGC TGC TCC CCG ACC CTG
 M N Q A M A T P L S L S C C S P T L
 ACC AGG TCC ACC CTG TTC TTC ACC AAG ACC TTC CCG TTC TCC CGC TCC TTC TCC
 T R S T L F F T K T F P F S R S F S
 ACC CCG CTG CCG CTG TCC ACC AAG ACC CTG ATC TCC CTG TCC CCG CCG CAC AGG
 T P L P L S T K T L I S L S P P H R
 ACC TTC GCC GTG AGG GCT GAG TCC CAG AAC GCC GCG GAC CCG GCC AGG CAG TAC
 E S Q N G A D P A R Q Y D F D L F T
 GAC TTC GAC CTG TTC ACC ATC GGC
 I G T F A V *R A

I. PHP9053 - Ubi::NLS-GFPm-MALS

The nuclear localization signal from simian virus 40 (SV40) was fused to the N terminus of GFPm. In order to retain the protein in the nucleus the molecular weight of NLS-GFPm was increased by making a carboxy terminal addition of the large unrelated protein maize acetolactate synthase (ALS) (lower case bases show linker sequence). Kalderon, D., Robers, B., Richardson, W., and Smith A., "A short amino acid sequence able to specify nuclear location", *Cell* 39: 499-509 (1984).

CCG CCC AAG AAG AAG CGC AAG GTG ccc ATG ...
 P P K K K R K V P M
 AAG atc cac ATG
 K I H M

J. Ubi::GFPm-HRGP

A carboxy terminal fusion of a portion of the maize HRGP coding sequence to GFPm was used to anchor GFP to the cell wall. Sequence encoding amino acids 177 to 328 was used. Stiefel, V., Ruiz-Avila, L., Raz R., Valles M., Gomez J., Pages M., Martinez-Izquierdo J., Ludevid M., Landale J., Nelson T., and Puigdomenech P., "Expression of a maize cell wall hydroxyproline-rich glycoprotein gene in early leaf and root vascular differentiation", *Plant Cell* 2: 785-793 (1990).

TAC ACT CCA AGC CCT AAG CCA CCG GCT ACC AAG CCT CCC ACG
 Y T P S P K P P A T K P P T

5 CCC AAG CCG ACC CCG CCA ACG TAC ACC CCT TCG CCA AAG CCT
 P K P T P P T Y T P S P K P
 CCG ACA CCC AAG CCG ACC CCG CCG ACG TAC ACC CCT TCT CCC
 P T P K P T P P T Y T P S P
 10 AAG CCT CCG ACG CCC AAG CCG ACC CCG CCG ACG TAC ACT CCA
 K P P T P K P T P P T Y T P
 AGC CCC AAG CCT CCC ACA CAC CCG ACG CCC AAG CCG ACC CCA
 S P K P P T H P T P K P T P
 CCG ACG TAC ACC CCT TCC CCA AAG CCT CCG ACG CCC AAG CCG
 P T Y T P S P K P P T P K P
 ACC CCA CCG ACG TAC ACC CCT TCC CCA AAG CCT CCG ACA CCC
 T P P T Y T P S P K P P T P
 15 AAG CCG ACC CCA CCG ACG TAC ACC CCT TCC CCA AAG CCT CCG
 K P T P P T Y T P S P K P P
 ACA CCC AAG CCG ACC CCA CCG ACG TAC ACT CCC ACA CCG AAG
 T P K P T P P T Y T P T P K
 CCG CCG GCC ACC AAG CCG CCC ACC TAC ACT CCG ACG CCG CCG
 P P A T K P P T Y T P T P P
 20 GTG TCT CAC ACC CCC AGC CCG CCG CCA CCA TAC TAC
 V S H T P S P P P P Y Y

K. Ubi::GFPm-SOD

25 Superoxide dismutase coding sequence fused to
 GFPm. Alcendor D., Chapman G., and Beaman B.,
 "Isolation, sequencing and expression of the
 superoxide dismutase-encoding gene (sod) of *Nocardia*
asteroides strain GUH-2", *Gene* 164: 143-147 (1995).

30 GTG GCT GAG TAC ACG CTG CCG GAT CTG GAT TAC GAC TAC AGC
 V A E Y T L P D L D Y D Y S
 GCC TGG AAC CCC ACA TCT CCG GGC AGA TCA ACG AGC TGA CAC
 A L E P H I S G Q I N E L H
 CAT TCC AAG CAC CAC GCC GCC TAC GTC GCC GGT GCC AAC ACG
 H S K H H A A Y V A G A N T
 35 GCA CTG GAG AAG CTG GAA GCC GCC COT GAG GCC GGC GAT CAC
 A L E K L E A A R E A G D H
 AGC GCG ATC TTC CTG CAC GAG AAG AAC CTC GCG TTC CAC CTC
 S A I F L H E K N L A F H L
 GGC GGA CAC GTC AAC CAC TCC ATC TGG TGG AAG AAC CTG TCC
 40 G G H V M H S I W W K N L S
 CCC AAC GGT GGC GAC AAG CCG GTC GGC GAG CTG GCC GCG GCC
 P N G G D K P V G E L A A A
 ATC GAC GAC CAG TTC GGT TCG TTC GAC AAG TTC CCG GCG CAG
 I D D Q F G S F D K F R A Q
 45 TTC ACC GCC GCG GCC AAC GGC CTG CAG GGC TCG GGC TGG GCG
 F T A A A N G L Q G S G W A
 GTG CTC GGT TAC GAC ACC CTC GGC CAG AAG CTG CTG ACC TTC
 V L G Y D T L G Q K L L T P
 CAG CTC TAC GAC CAG CAG GCC AAC GTG CCG CTG GGC ATC ATC
 50 Q L Y D Q Q A M V P L G I I
 CCG CTG CTC CAG GTC GAC ATG TGG GAG CAC GCC TTC TAC CTG
 P L L Q V D M W E H A F Y L
 CAG TAC AAG AAC GTC AAG GCC GAC TAC GTG ACC GCG TTC TGG
 Q Y K N V K A D Y V T A F W

AAC GTC GTC AAC TGG GCC GAC GTG CAG GAC CGC TTC GGC AAG
N V V N W A D V Q D R F G K
GCC GTC AAC CAG GGC AAG GGC CTT ATC TTC GGG
A V N Q G K G L I F G

5

9. Detection of GFP in Transformed Plant Cells

GFP is detected in transformed plant cells by conventional spectrophotometric methods. The transformed cells or tissue are screened for the presence of GFP protein by means of the illuminating the cells with UV to blue light and screening for the presence of green fluorescence.

Compound and dissecting microscopes were fitted with appropriate filter combinations for fluorescent protein excitation. Illumination with UV-blue light (exciting around the absorption maximum of 395 nm or around the minor peak at approximately 475 nm for GFP; around 480-490 nm for the red-shifted version and around 380 nm for the blue fluorescent protein) is required for visualization. A hand-held lamp for benchtop work also permits good visualization. Cut-off filters or bandpass filters between the fluorescing tissue and the viewer (i.e. around the intermediate objective or the eyepieces of the microscope, or hand-held in front of the eyes if working on the benchtop) greatly reduced background autofluorescence from the tissue. For GFP and the red-shifted GFP this cut-off or bandpass filter permitted light between 500-550 nm to reach the viewer. For the blue fluorescent protein, this filter permitted the passage of 420 to 450 nm light. Useful filters and wavelengths are not restricted to those described above, and further generic description of the optical characteristics of this system are available. Heim et al., *Currently Biology* 6: 178-182 (1996).

10. Foreign Protein Genes and Agronomic Genes

With transgenic plants according to the present invention, a foreign protein can be produced in commercial quantities. Thus, the selection and propagation techniques described above yield a plurality of transgenic plants which are harvested in a conventional manner, and a foreign protein then is extracted from a tissue of interest or from total biomass. Protein extraction from plant biomass can be accomplished by known methods which are discussed, for example, by Heney and Orr, *Anal. Biochem.* 114: 92-6 (1981).

According to a preferred embodiment, the transgenic plant provided for commercial production of foreign protein is maize. In another preferred embodiment, the biomass of interest is seed. For the relatively small number of transgenic plants that show higher levels of expression, a genetic map can be generated, primarily via conventional RFLP and PCR analysis, which identifies the approximate chromosomal location of the integrated DNA molecule. For exemplary methodologies in this regard, see Glick and Thompson, *METHODS IN PLANT MOLECULAR BIOLOGY AND BIOTECHNOLOGY* 269-284 (CRC Press, 1993). Map information concerning chromosomal location is useful for proprietary protection of a subject transgenic plant. If unauthorized propagation is undertaken and crosses made with other germplasm, the map of the integration region can be compared to similar maps for suspect plants, to determine if the latter have a common parentage with the subject plant. Map comparisons would involve hybridizations, RFLP, PCR and sequencing, all of which are conventional techniques.

Likewise, by means of the present invention, agronomic genes can be expressed in transformed plants. More particularly, plants can be genetically engineered to express various phenotypes of agronomic interest. The

genes implicated in this regard include, but are not limited to, those categorized below.

5 1. Genes That Confer Resistance To Pests or Disease And That Encode:

10 (A) Plant disease resistance genes. Plant defenses are often activated by specific interaction between the product of a disease resistance gene (R) in the plant and the product of a corresponding avirulence (Avr) gene in the pathogen. A plant variety can be transformed with cloned resistance gene to engineer plants that are resistant to specific pathogen strains. See, for example Jones et al., *Science* 266: 789 (1994) (cloning of the tomato Cf-9 gene for resistance to *Cladosporium fulvum*); Martin et al., *Science* 262: 1432 (1993) (tomato Pto gene for resistance to *Pseudomonas syringae* pv. tomato encodes a protein kinase); Mindrinos et al., *Cell* 78: 1089 (1994) (*Arabidopsis* RSP2 gene for resistance to *Pseudomonas syringae*).

25 (B) A *Bacillus thuringiensis* protein, a derivative thereof or a synthetic polypeptide modeled thereon. See, for example, Geiser et al., *Gene* 48: 109 (1986), who disclose the cloning and nucleotide sequence of a Bt δ -endotoxin gene. Moreover, DNA molecules encoding δ -endotoxin genes can be purchased from American Type Culture Collection (Rockville, MD), under ATCC accession Nos. 40098, 67136, 31995 and 31998.

30 (C) A lectin. See, for example, the disclosure by Van Damme et al., *Plant Molec. Biol.* 24: 825 (1994), who disclose the nucleotide sequences of several *Clivia miniata* mannose-binding lectin genes.

35 (D) A vitamin-binding protein such as avidin. See U.S. patent application serial No. 07/911,864, the contents of which are hereby incorporated by reference.

The application teaches the use of avidin and avidin homologues as larvicides against insect pests.

5 (E) An enzyme inhibitor, for example, a protease inhibitor or an amylase inhibitor. See, for example, Abe et al., *J. Biol. Chem.* 262: 16793 (1987) (nucleotide sequence of rice cysteine proteinase inhibitor), Huub et al., *Plant Molec. Biol.* 21: 985 (1993) (nucleotide sequence of cDNA encoding tobacco proteinase inhibitor I), and Sumitani et al., *Biosci. Biotech. Biochem.* 57: 1243 (1993) (nucleotide sequence of *Streptomyces nitrosporeus* α -amylase inhibitor).

15 (F) An insect-specific hormone or pheromone such as an ecdysteroid and juvenile hormone, a variant thereof, a mimetic based thereon, or an antagonist or agonist thereof. See, for example, the disclosure by Hammock et al., *Nature* 344: 458 (1990), of baculovirus expression of cloned juvenile hormone esterase, an inactivator of juvenile hormone.

25 (G) An insect-specific peptide or neuropeptide which, upon expression, disrupts the physiology of the affected pest. For example, see the disclosures of Regan, *J. Biol. Chem.* 269: 9 (1994) (expression cloning yields DNA coding for insect diuretic hormone receptor), and Pratt et al., *Biochem. Biophys. Res. Comm.* 163: 1243 (1989) (an allostatin is identified in *Diploptera punctata*). See also U.S. patent No. 5,266,317 to Tomalski et al., who disclose genes encoding insect-specific, paralytic neurotoxins.

35 (H) An insect-specific venom produced in nature by a snake, a wasp, etc. For example, see Pang et al., *Gene* 116: 165 (1992), for disclosure of heterologous expression in plants of a gene coding for a scorpion insectotoxic peptide.

(I) An enzyme responsible for an hyperaccumulation of a monerpene, a sesquiterpene, a steroid, hydroxamic acid, a phenylpropanoid derivative or another non-protein molecule with insecticidal activity.

5

(J) An enzyme involved in the modification, including the post-translational modification, of a biologically active molecule; for example, a glycolytic enzyme, a proteolytic enzyme, a lipolytic enzyme, a nuclease, a cyclase, a transaminase, an esterase, a hydrolase, a phosphatase, a kinase, a phosphorylase, a polymerase, an elastase, a chitinase and a glucanase, whether natural or synthetic. See PCT application WO 93/02197 in the name of Scott et al., which discloses the nucleotide sequence of a callase gene. DNA molecules which contain chitinase-encoding sequences can be obtained, for example, from the ATCC under accession Nos. 39637 and 67152. See also Kramer et al., *Insect Biochem. Molec. Biol.* 23: 691 (1993), who teach the nucleotide sequence of a cDNA encoding tobacco hookworm chitinase, and Kawalleck et al., *Plant Molec. Biol.* 21: 673 (1993), who provide the nucleotide sequence of the parsley ubi4-2 polyubiquitin gene.

10

15

20

25

(K) A molecule that stimulates signal transduction. For example, see the disclosure by Botella et al., *Plant Molec. Biol.* 24: 757 (1994), of nucleotide sequences for mung bean calmodulin cDNA clones, and Griess et al., *Plant Physiol.* 104: 1467 (1994), who provide the nucleotide sequence of a maize calmodulin cDNA clone.

30

(L) A hydrophobic moment peptide. See U.S. patent applications serial No. 08/168,809 (disclosure of peptide derivatives of Tachyplesin which inhibit fungal plant pathogens) and serial No. 08/179,632 (teaches synthetic antimicrobial peptides that confer disease resistance),

35

the respective contents of which are hereby incorporated by reference.

5 (M) A membrane permease, a channel former or a channel blocker. For example, see the disclosure by Jaynes et al., *Plant Sci.* 89: 43 (1993), of heterologous expression of a cecropin- β lytic peptide analog to render transgenic tobacco plants resistant to *Pseudomonas solanacearum*.

10 (N) A viral-invasive protein or a complex toxin derived therefrom. For example, the accumulation of viral coat proteins in transformed plant cells imparts resistance to viral infection and/or disease development effected by the virus from which the coat protein gene is
15 derived, as well as by related viruses. See Beachy et al., *Ann. Rev. Phytopathol.* 28: 451 (1990). Coat protein-mediated resistance has been conferred upon transformed plants against alfalfa mosaic virus, cucumber
20 mosaic virus, tobacco streak virus, potato virus X, potato virus Y, tobacco etch virus, tobacco rattle virus and tobacco mosaic virus. *Id.*

25 (O) An insect-specific antibody or an immunotoxin derived therefrom. Thus, an antibody targeted to a critical metabolic function in the insect gut would inactivate an affected enzyme, killing the insect. Cf. Taylor et al., Abstract #497, SEVENTH INT'L SYMPOSIUM ON MOLECULAR PLANT-MICROBE INTERACTIONS (1994) (enzymatic
30 inactivation in transgenic tobacco via production of single-chain antibody fragments).

35 (P) A virus-specific antibody. See, for example, Tavladoraki et al., *Nature* 366: 469 (1993), who show that transgenic plants expressing recombinant antibody genes are protected from virus attack.

(Q) A developmental-arrestive protein produced in nature by a pathogen or a parasite. Thus, fungal endo α -1,4-D-polygalacturonases facilitate fungal colonization and plant nutrient release by solubilizing plant cell wall homo- α -1,4-D-galacturonase. See Lamb et al., *Bio/Technology* 10: 1436 (1992). The cloning and characterization of a gene which encodes a bean endopolygalacturonase-inhibiting protein is described by Toubart et al., *Plant J.* 2: 367 (1992).

(R) A developmental-arrestive protein produced in nature by a plant. For example, Logemann et al., *Bio/Technology* 10: 305 (1992), have shown that transgenic plants expressing the barley ribosome-inactivating gene have an increased resistance to fungal disease.

2. Genes That Confer Resistance To A Herbicide, For Example:

(A) A herbicide that inhibits the growing point or meristem, such as an imidazalinone or a sulfonylurea. Exemplary genes in this category code for mutant ALS and AHAS enzyme as described, for example, by Lee et al., *EMBO J.* 7: 1241 (1988), and Miki et al., *Theor. Appl. Genet.* 80: 449 (1990), respectively.

(B) Glyphosate (resistance imparted by mutant EPSP synthase and *aroA* genes, respectively) and other phosphono compounds such as glufosinate (PAT and *bar* genes), and pyridinoxy or phenoxy proprionic acids and cyclohexones (ACCase inhibitor-encoding genes). See, for example, U.S. patent No. 4,940,835 to Shah et al., which discloses the nucleotide sequence of a form of EPSP which can confer glyphosate resistance. A DNA molecule encoding a mutant *aroA* gene can be obtained under ATCC accession No. 39256, and the nucleotide sequence of the mutant gene is disclosed in U.S. patent No. 4,769,061 to Comai. European patent application No. 0 333 033 to

Kumada et al. and U.S. patent No. 4,975,374 to Goodman et al. disclose nucleotide sequences of glutamine synthetase genes which confer resistance to herbicides such as L-phosphinothricin. The nucleotide sequence of a phosphinothricin-acetyl-transferase gene is provided in European application No. 0 242 246 to Leemans et al. De Greef et al., *Bio/Technology* 7: 61 (1989), describe the production of transgenic plants that express chimeric bar genes coding for phosphinothricin acetyl transferase activity. Exemplary of genes conferring resistance to phenoxy proprionic acids and cyclohexones, such as sethoxydim and haloxyfop, are the Acc1-S1, Acc1-S2 and Acc1-S3 genes described by Marshall et al., *Theor. Appl. Genet.* 83: 435 (1992).

(C) A herbicide that inhibits photosynthesis, such as a triazine (*psbA* and *gs+* genes) and a benzonitrile (nitrilase gene). Przibilla et al., *Plant Cell* 3: 169 (1991), describe the transformation of *Chlamydomonas* with plasmids encoding mutant *psbA* genes. Nucleotide sequences for nitrilase genes are disclosed in U.S. patent No. 4,810,648 to Stalker, and DNA molecules containing these genes are available under ATCC accession Nos. 53435, 67441 and 67442. Cloning and expression of DNA coding for a glutathione S-transferase is described by Hayes et al., *Biochem. J.* 285: 173 (1992).

3. Genes That Confer Or Contribute To A Value-Added Trait, Such As:

(A) Modified fatty acid metabolism, for example, by transforming a plant with an antisense gene of stearoyl-ACP desaturase to increase stearic acid content of the plant. See Knultzon et al., *Proc. Nat'l Acad Sci. USA* 89: 2624 (1992).

(B) Decreased phytate content

(1) Introduction of a phytase-encoding gene would enhance breakdown of phytate, adding more free phosphate to the transformed plant. For example, see Van Hartingsveldt et al., *Gene* 127: 87 (1993), for a disclosure of the nucleotide sequence of an *Aspergillus niger* phytase gene.

(2) A gene could be introduced that reduces phytate content. In maize, this, for example, could be accomplished, by cloning and then re-introducing DNA associated with the single allele which is responsible for maize mutants characterized by low levels of phytic acid. See Raboy et al., *Maydica* 35: 383 (1990).

(C) Modified carbohydrate composition effected, for example, by transforming plants with a gene coding for an enzyme that alters the branching pattern of starch. See Shiroza et al., *J. Bacteriol.* 170: 810 (1988) (nucleotide sequence of *Streptococcus mutans* fructosyltransferase gene), Steinmetz et al., *Mol. Gen. Genet.* 200: 220 (1985) (nucleotide sequence of *Bacillus subtilis* levansucrase gene), Pen et al., *Bio/Technology* 10: 292 (1992) (production of transgenic plants that express *Bacillus licheniformis* α -amylase), Elliot et al., *Plant Molec. Biol.* 21: 515 (1993) (nucleotide sequences of tomato invertase genes), Søgaard et al., *J. Biol. Chem.* 268: 22480 (1993) (site-directed mutagenesis of barley amylase gene), and Fisher et al., *Plant Physiol.* 102: 1045 (1993) (maize endosperm starch branching enzyme II).

Example 1

Construction of Genes and Vectors

5 The GFPm nucleotide sequence was derived from
a back translation of the protein sequence using maize
preferred codons. Sequence analysis was performed using
the Wisconsin Sequence Analysis Package from Genetics
10 Computer Group, Madison, WI. The nucleotide sequence was
assembled from a series of synthetic oligonucleotides.
Cloning sites include a 5-prime flanking *Bam*HI
restriction site, an *Afl*III site at the start codon, a 3-
prime flanking *Hpa*I site or a *Bgl*II site converting the
stop codon to an isoleucine (Fig. 1).

15 Amino terminal fusions to GFPm were created using
synthetic oligonucleotides encoding the signal sequence
flanked by a 5-prime *Bam*HI or *Bgl*II restriction site and
3-prime *Nco*I site. The synthetic signal sequences were
cloned into PHP7921 (Fig. 4) using the 5-prime flanking
20 *Bam*HI and *Afl*III restriction sites of GFPm to create in-
frame fusions. Sequences cloned in this manner include
the maize chloroplast signal sequence, barley alpha-
amylase signal sequence, pea glutathione reductase signal
sequence, and SV40 nuclear localization signal. The
25 sporamin vacuole signal sequence was synthesized with a
5-prime *Bsp*EI and 3-prime *Bam*HI for insertion between the
BAA pre-sequence and GFPm coding sequence of pHP8144
(Fig. 5). The CPN60 promoter site, created by site-
directed mutagenesis, was used with the GFPm *Afl*III site
30 to make the fusion sequence. Each of plasmids PHP7921
and PHP8144 are based on pUC18.

35 Carboxy terminal sequences were synthesized with
a 5-prime *Bgl*II and 3-prime *Hpa*I site, and cloned into
PHP7921 or pHP8144 using the 3-prime *Bfl*II and *Hpa*I sites
of GFPm. The barley lectin vacuole target, HDEL ER
retention signal, SKL peroxisome signal, ALS coding
sequence and HRGP coding sequence fusions were made using
this strategy.

TABLE 1
PLASMIDS

PHP Number	Target Crop	Promoter	Targeting				Other 5' Sequences	Structural Gene	3' Sequence	Comments
			5' Sequence	3' Sequence	Localized To:					
7956	maize	4xERE	--	--	Cytoplasm		Omega Prime (O') Adhl intron I	GFPm	pinII	Estrogen-inducible expression
7667	maize	ubiquitin	--	--	Cytoplasm		--	GFP (native)	nos	Jellyfish structural gene
7921	maize	ubiquitin	--	--	Cytoplasm		--	GFPm	pinII	Cytoplasmic expression
8489	maize	ubiquitin	--	--	Cytoplasm		--	GFPm	pinII	MAR flanking cassette
8490	maize	ubiquitin	--	--	Cytoplasm		--	GFPm-S65T	pinII	MAR flanking cassette, red-shift
8080	maize	ubiquitin	CTP	--	chloroplast		--	GFPm	pinII	Chloroplast signal sequence, cab-m7 from maize

8087	maize	CPN60	CPN60	--	mitochondria a	--	GFPm	pinII	Targeted to mitochondria, signal from maize chaperonin 60
8144	maize	ubiquitin	BAA	--	secreted	--	GFPm	pinII	Secreted GFP; signal from barley alpha amylase 1
8757	maize	ubiquitin	BAA	BLVT	vacuole	--	GFPm	pinII	Barley lectin vacuole signal seq. on carboxy end of BAA- GFPm
8758	maize	ubiquitin	BAA	HDEL	endoplasmic reticulum	--	GFPm	pinII	HDEL endoplasmic reticulum signal from maize b-70
8759	maize	ubiquitin	BAA	--	vacuole	SVT	GFPm	pinII	Sweet potato sporamin vacuole signal sequence

8882	maize	ubiquitin	--	SKL	peroxisome	--	GFPm	pinII	Consensus peroxisomal signal sequence at carboxy terminus
8986	maize	ubiquitin	T7 TAG-NLS-	--	nucleus	--	GFPm	pinII	Nuclear localization sequence
9053	maize	ubiquitin	T7 TAG-NLS-	--	nucleus	--	MALS-GFPm	pinII	Nuclear localization sequence; protein fusion to prevent escape
9057	maize	ubiquitin	--	--	nucleus	--	MALS-GFPm	pinII	Fusion protein to test effect on GFP as reporter
9058	maize	ubiquitin	--	--	cytoplasm	--	PATm-GFPm	pinII	PAT-GFP fusion to test efficacy

8744	maize	ubiquitin	--	--	nucleus	--	GFPm-ZmRAD51	pinII	Protein fusion: for studies on Rad51 expression/function
8081	maize	ubiquitin	--	--	cytoplasm	FRT	GFPm	pinII	To verify that FRT sequence does not impair expression
8674	maize	2x35s	--	--	cytoplasm	O' Adh1 intron	FRT-bar-FRT-GFPm	pinII	Select on bialaphos, then excise bar and activate GFP
8911	maize	ubiquitin / 2x35s	--	adh1 intron 1	cytoplasm	Ds	GFPm bar	pinII pinII	ubi-Ds-2x35S: :bar::pin-Ds-GFPm::pinII
8088	maize	ubiquitin	--	--	cytoplasm	--	GFPm-S65T	pinII	Red-shifted GFPm
8138	maize	ubiquitin	--	--	cytoplasm	--	GFPm-Red2	pinII	Red-shifted version #2 F64M, S65G, Q69L

8175	maize	ubiquitin	--	--	cytoplasm	--	Blue FP	pinII	GFPm-modified Y66H
8007	maize	2x35s ubiquitin					adh1 intronI::bar ubi intron::FLP	pinII pinII	Tandem vector for FLP excision of FRT to activate GFPm
8011	sunflo wer	2x35s ALS	--	--	cytoplasm	0'	GFPm NPT-II	pinII pinII	Binary
8491	sunflo wer	2x35s ALS	--	--	cytoplasm	0'	GFPm-S65T NPT-II	pinII pinII	Binary with red-shifted GFP

The ubiquitin promoter is described in Christensen et al., *Plant Mol. Biol.* 18: 675-689 (1992). The 4xERE promoter is described in Klein-Hitpass et al., *Cell* 46: 1053-1061 (1986). The CPN60; 2x35S and ALS promoters are described in Close P. Ph.D. Dissertation, Iowa State University; Gardner et al., *Nucl. Acid Res.* 9: 2871-2888 (1981) and PHI Patent Application 08/409,297, respectively. The Omega prime (O') 5-prime sequence is described by Gallie et al., *Nucl. Acids Res.* 15: 3257-3273 (1987) while the Adh 1 intron 1 is described by Dennis et al., *Nucl. Acids Res.* 12: 3983-3990 (1984). The 3-prime sequence pinII is described in An et al., *Plant Cell* 1: 115-122 (1989). Structural genes used in the construction of the vectors include the GFP gene (native jellyfish) described by Prasher et al., *Gene* 111: 229-233 (1992); the maize ALS gene described by Fromm et al., *Bio/Technol.* 8: 883 (1990) and the maize HPRG described by Steifel et al., *Plant Mol. Biol.* 11: 483-493 (1988). The FLP/FRT system contained in plasmids PHP8674 and PHP8007 is described in O'Gorman et al., *Science* 251: 1351-1355 (1991) and Jayaram M., *Proc. Natl. Acad. Sci. USA* 82: 5875-5879 (1985). The Ac/Ds system is described by Muller-Neumann et al., *Mol. Gen. Genet.* 198: 19-24 (1984).

For *Agrobacterium*-mediated transformation of sunflower, the backbone of PHP8011 is a bin 19-based vector. Beven, *Nucl. Acids Res.* 12: 8711-8721 (1984). The NPT-II cassette was removed from bin 19 and the following cloned between the left and right border sequences; the double 35S promoter, omega', the GFPm coding sequence and the pinII 3' region (all described above) were cloned downstream of the left border sequence, followed in a head-to-tail orientation by the BnALS3 promoter, the NPT-II coding sequence and the pinII 3' region. Beven, "Binary *Agrobacterium* Vectors for Plant Transformation", *Nucl. Acids Res.* 12: 8711-8721 (1984).

GFP_r, BFP and GFPs (soluble GFP) were constructed by means of oligonucleotide-directed mutagenesis of the GFP_m nucleotide sequence shown in Figure 1 as described by Ausubel, *supra*. With regard to GFP_r, TCC was changed to ACG resulting in the S65T substitution in the GFP amino acid sequence. With regard to BFP, TAC was changed to CAC, resulting in the Y66H substitution in the GFP amino acid sequence.

With regard to GFPs, the following sequence changes in GFP_m were made. TCC was changed to AGC, resulting in a F99S substitution in the GFP amino acid sequence. The oligonucleotide used to introduce this mutation also caused a silent mutation at R96 (AGC changed to CCG), thereby adding a *Bsr*BI restriction site. ATG was changed to ACC, resulting in an M153T substitution in the GFP amino acid sequence. GTG was changed to GCC, resulting in a V163A substitution in the GFP amino acid sequence. A single oligonucleotide was used to change the amino acids at 153 and 163. This oligonucleotide also caused a silent mutation at A154 (GCC changed to GCG), which added a *Sac*II restriction site. As reported by Cramer *et al.*, *supra*, these three amino acid changes did not alter the absorption/emission characteristics of GFP. These three amino acid substitutions, however, increased the solubility of GFP. Maize cells transformed with the GFPs gene fluoresce in blue to UV light based on transient assays.

Example 2

Maize Transformation with GFP Screening or Bialaphos Selection

Maize Hi-II immature embryos were co-transformed via microprojectile bombardment with a mixture of PHP7814 (ubi::PAT::pinII) and PHP8409. Plasmid PHP7814 carries the PAT gene encoding resistance to bialaphos operably linked to the ubiquitin promoter and the terminator from the pinII gene. Standard CaCl₂/spermidine precipitation of DNA onto tungsten microprojectiles was used to prepare the DNA-particles. Prior to bombardment, the embryos of approximately 1.5-2.0 mm in length were cultured on 560P medium including N6 salts, Erikkson's vitamins, 0.69 mg/L proline, 2 mg/L 2,4-D, and 3% sucrose. After 4-5 days of incubation in the dark at 28°C, embryos were removed from 560P medium and cultured, coleorhizal end up, on 560L medium which is equivalent to 560P except that it contains 12% sucrose. Embryos were allowed to acclimate to this medium for 3 h prior to transformation.

Embryos were transformed using the PDS-1000 Helium Gun from Bio-Rad at one shot per sample using 650 PSI rupture disks. On average approximately 0.0067 µg of DNA was delivered per shot. For each treatment (pat gene with bialaphos selection or GFP gene with screening for GFP), 447 embryos were used. For the bialaphos-selection treatment; following bombardment, the embryos were maintained on 560L medium for 48 h before transfer to 560R selection medium containing N6 salts, Erikkson's vitamins, 2 mg/L 2,4-D, 3% sucrose and 3mg/L bialaphos. Plates were maintained at 28°C in the dark and were observed for colony recovery. Transfer of colonies to fresh medium occurred every two weeks. Transgenic colony recovery under bialaphos

selection was accomplished by picking the healthy, growing transgenic calli from amongst the inhibited background of non-transgenic tissue as the material was progressively transferred to fresh media. For the GFP screening treatment, 560P media was used after bombardment and the two-day recovery on 560L. Calli were transferred to fresh non-selective 560P medium every two weeks. Screening for GFP expression was carried out at each transfer using a Xenon and/or Mercury light source with the appropriate filters for GFP visualization.

Table 2

Recovery of Transgenic Events in Maize Hi-II Callus Using Either GFP Visualization or Bialaphos Selection

Selection Method	Number of Events Recovered	
	Initial Events	Events Capable of Regeneration
Bialaphos selection	16	12
Screened using GFP	9	8

Transgenic events were recovered from each treatment. The largest number of transgenic events were consistently obtained by means of bialaphos selection. However, the frequency of recovery of transgenic events with GFP screening was approximately one-half the rate of recovery with bialaphos selection which was surprisingly high considering that GFP screening does not rely on selection of transgenic events in the presence of a cellular toxin.

Once GFP expressing colonies were identified they were monitored regularly for new growth and expression using the Xenon light source. Plant cells containing GFP were regenerated by transferring the callus to 288 medium containing MS salts, 1 mg/L IAA,

0.5 mg/L zeatin and 4% sucrose. The callus was placed in the light. As plantlets developed they were transferred to tubes containing 272K, hormone-free MS medium and 3% sucrose. The percentage of green
5 fluorescent colonies that regenerated into whole plants was determined.

Leaf samples were collected from T0 transgenic plants from events recovered by means of bialaphos selection, or for events recovered by means of GFP
10 visualization (non-selected). Genomic DNA was extracted from all leaf samples and Southern analysis was performed using the PAT and GFP genes as hybridization probes. Hybridization for the PAT gene and GFP gene were performed sequentially on the same
15 blots, by stripping and reprobing for the other transgene. Bands hybridizing for the PAT gene and for the GFP gene were observed in all 8 GFP-screened events tested and in all 12 of the bialaphos selected events tested. Copy numbers and integration patterns for both
20 treatments appeared similar, and were consistent with previous experiments by the inventors using bialaphos selection of callus. Copy numbers ranged from low (single or two copies for each gene) to numerous (i.e., >5 copies per gene). Integration events were
25 predominantly at single integration sites but in some cases integration at multiple locations in the genome was observed.

Bialaphos is the most efficient chemical selective agent for maize. Consequently, the high
30 frequency with which transgenic events were obtained with GFP screening, compared to bialaphos selection, was unexpected. An advantage of GFP screening over bialaphos selection is that non-transformed tissue is not killed. Accordingly, transgenic sectors containing
35 the GFP gene develop in a healthier environment for growth and development.

Example 3**The GFP Marker in Maize Cytoplasm, Chloroplast and Mitochondria**

5 Maize A188 x B73-derived cells were transformed
with plasmids PHP8080, PHP7921 and PHP8087. Each of
these plasmids is described in Example 1 and Table 1.
Plasmid PHP8080 carries the GFPm gene operably linked
to the ubiquitin promoter and a chloroplast signal
10 sequence. GFP is directed to the chloroplasts of cells
transformed with PHP8080. Plasmid PHP7921 carries the
GFPm gene operably linked to ubiquitin promoter. GFP
remains in the cytoplasm of cells transformed with
pPGP7921. Plasmid PHP8087 carries the GFPm gene
15 operably lined to the CPN60 promoter and CPN60 signal
sequence. GFP is directed to mitochondria of cells
transformed with pPGP8087.

 The cells were transformed by culturing maize
embryos approximately 1.5 to 2 mm in length onto 560P
20 medium containing N6 salts, Eriksson's vitamins, 0.69
g/L Proline, 2 mg/L 2,4-D and 3% sucrose. After 4-5
days of incubation in the dark at 28°C, embryos were
removed from 560P medium and cultured, coleorhizal end
up, onto 560L medium which is equivalent to 560P but
25 contains 12% sucrose. Embryos were allowed to
acclimate to this medium for 3 h prior to
transformation. Embryos were transformed using the
PDS-1000 Helium Gun from Bio-Rad at one shot per sample
using 650PSI rupture disks. DNA delivered per shot
30 averaged at 0.0667 µg. Following bombardment, all
embryos were maintained on 560L medium for 48 hours
before transfer to 3% sucrose and 3 mg/L bialaphos.
Plates were maintained at 28°C in the dark and were
observed for colony recovery with transfers to fresh
35 medium occurring every two weeks. Transgenic colony
recovery was noted initially as growing callus tissue
with a healthy phenotype on selection. Screening for

GFP expression was completed with each transfer using a Xenon light source with UV filter attachments.

GFP localization was confirmed by analysis of TO callus. Samples of calli from the targeted GFP transformations were fixed in FAA and then examined using an inverted microscopes with UV filters to visualize GFP. For each targeted construct GFP expression was localized to the specified organelle.

Once GFP expressing colonies were identified they were monitored regularly for new growth and expression using the Xenon light source. Plant cells containing GFP were regenerated by transferring the callus to 288 medium containing MS salts, 1 mg/L IAA, 0.5 mg/L zeatin and 4% sucrose. The callus was placed in the light. As plantlets developed they were transferred to tubes containing 272K, hormone-free MS medium and 3% sucrose. The percentage of green fluorescent colonies that regenerated into whole plants was determined.

Regeneration of plants from callus transformed with PHP8080, PHP7921, and PHP8087 appeared to vary between constructs; plants were successfully regenerated from independent transformed calli at respective frequencies of 35, 54 and 86%. Transformation was confirmed by Southern blot analyses using both the BAR and GFPm genes as hybridization probes. GFP localization in various subcellular compartments was confirmed in stable transgenic maize cells using epifluorescent microscopy and image enhancement software.

Table 3

**Regeneration Frequency Of Maize Transformed With GFP
Directed to Different Subcellular Compartments**

Chloroplast					
Exp #	Total Colonies	Green	Regenerate	Percentage	Ave. Plant Event
..4.01	32	23	4	17	6
..4.03	0	-	-	-	-
..4.06	13	11	8	73	3
Totals	45	34	12	35*	
Cytoplasm					
Exp #	Total Colonies	Green	Regenerate	Percentage	Ave. Plants Event
..4.01	7	6	2	33	4
..4.03	7	5	4	80	5
..4.06	8	7	6	86	4
..4.08	9	9	3	33	2
..4.11	3	3	1	33	4
Totals	34	30	16	54*	
Mitochondrial					
Exp #	Total Colonies	Green	Regenerate	Percentage	Ave. Plants Event
..4.03	5	3	1	33	9
..4.06	13	11	11	100	4
Totals	18	14	12	86*	
* This is not a mean regeneration frequency, but instead is calculated from the totals for all treatments.					

Transformed plants containing GFP in the mitochondrial subcellular compartment grew more vigorously than plants containing GFP in the cytoplasm or chloroplast. The transformed plants containing GFP in the mitochondria were equivalent to the controls in terms of growth rate and appearance.

In contrast, transformed plants containing GFP in the chloroplast were difficult to regenerate. This callus produced a preponderance of roots with few shoots. Almost all the plants containing GFP in the chloroplast that were regenerated and reached maturity clearly exhibited stress with narrow and somewhat flaccid leaves, reduced stature, yellow leaves (presumably caused by reduced chlorophyll levels) and male sterility.

Germination of T1 embryos have been tested in the following manner. Aliquots of seed from a number of independent transformants were imbibed and the embryos were excised and placed on hormone-free MS medium with 3% sucrose for germination (this in vitro test made it easier to visualize the seedling). All embryos were germinated in the dark and then scored for visual GFP expression using UV-blue excitation under the dissecting microscope.

For each event tested aliquots of 5 or 10 seeds/event were imbibed, excised and scored for germination and fluorescence. The germination frequency for excised progeny produced on plants transformed with PHP7921 and PHP8080 was good. The germination frequency for excised progeny produced on plants transformed with PHP8087 was good for 2 events (10/10), low for one event (2/10). No detectable germination was observed for 2 events. This lowered germination frequency could be due to mitochondrially-expressed GFP expression per se, but it could also be the result of event-to-event variation sometimes observed in transgenic lines.

Germination of a single event from plants transformed with PHP7921 was tested differently. Seed were screened using a UV-blue light source, and 10 seed with GFP-expressing seed (embryos and endosperm) were selected. Five of these seed were planted in soil, and in the other 5 the embryos were excised and germinated on medium. None of the seed in soil germinated, while

all excised embryos germinated, grew and the seedlings continued to fluoresce. For events where germination appears to be negatively impacted, this could be due to increased GFP concentrations during seed maturation and dry-down. High GFP concentrations and aggregation of the protein has been found to be toxic in *E. coli*. Cramer et al., *Nature Biotechnol.* 14: 315-319 (1995).

5

Table 4
T1 Progeny Analysis
Plant Germination and GFPm Expression

Plasmid	Event Tested	# of T1 Seed Tested	# of T1 Seed Germinated Per Event	# of T1 Plants Expressing GFPm
PHP8087: CPN60::mit-GFPm-PinII	1	10	0	0
PHP8087: CPN60::mit-GFPm-PinII	2	10	0	0
PHP8087: CPN60::mit-GFPm-PinII	3	10	10	1
PHP8087: CPN60::mit-GFPm-PinII	4	10	2	0
PHP8087: CPN60::mit-GFPm-PinII	5	10	10	2
PHP7921: Ubi::Ubi intron-GFPm-PinII	1	10	10	4
PHP8080: Ubi::Ubi intron-CTP-GFPm-PinII	1	5	5	0
PHP8080: Ubi::Ubi intron-CTP-GFPm-PinII	2	5	4	0
PHP8080: Ubi::Ubi intron-CTP-GFPm-PinII	3	5	4	0
PHP4810: Ubi::Ubi intron-BAR-PinII	1	5	5	0
*PHP7921: Ubi::Ubi intron-GFPm-PinII	2	10 5 germinated in vitro and 5 germinated in soil	5 Germination occurred only from in vitro cultures	5 Seed planted in soil also were GFPm +

* For all seed except those designated in the PHP7921 transformation, embryos were excised from imbibed seed and germinated on MS medium containing no hormones and 3% sucrose. All embryos were germinated in the dark to allow for visualization of GFPm expression. One event from PHP7921 was tested by planting one half of the seed in soil and one half with embryos excised from endosperm with culture in vitro. All seed planted in soil did not germinate. The germination rate for embryos excised from the endosperm and cultured in vitro was 100%. All plants expressed GFPm.

Example 4

Sunflower Transformation with GFP Screening, GUS Screening and Kanamycin Selection

5 Experiments were undertaken to test GFP, GUS and NPTII as marker genes for sunflower transformation. Two different kinds of meristem systems (split
10 meristem; intact meristem) and two different kinds of particle systems (FG, focusing gun; He, helium gun) were used. The screenable markers GFP and GUS were tested with the intact meristem system under conditions
15 involving no chemical selection but rather selection of transformants based on visual (GFP) or assayable (GUS) expression in transformed sectors. The NPTII marker gene was tested with the split meristem method using kanamycin for chemical selection of transformants and assays for NPTII expression to identify transformed
20 sectors. In some cases, the experiments were carried only so far as identifying transformed sectors while other experiments were carried out all the way through to test for seed transmission of the marker gene to progeny. Transmission to progeny was confirmed by
25 NPTII ELISA.

Regardless of the exact meristem system or type of particle gun used, the basic transformation protocol involved a combination of wounding by particle bombardment, followed by use of *Agrobacterium* for DNA
30 delivery, as described by Bidney et al., *Plant Mol. Biol.* 18: 301-313 (1992). The plasmids used for plant transformation in these experiments were PHP167 containing GUS as the screenable marker, PHP8011 containing GFP as the screenable marker as well as the
35 NPTII gene, and PHP762 containing NPTII as the selectable marker. Both PHP167 and PHP762 are binary vectors derived from pGA473. See An et al., *EMBO J.* 4: 277-284 (1985). The binary backbone for PHP8011 is bin 19. See Bevan, *NAR* 12: 8711-8721 (1984).

Procedures for preparation of *Agrobacterium* and preparation of particles for all of the experiments are described in Bidney et al., *supra*. The Pioneer sunflower line SMF3 was used in all experiments. See Burrus et al., *Plant Cell Rep.* 10: 161-166 (1991) for information concerning SMF3. The *Agrobacterium* strain EHA101 is described in Bidney et al. *supra* and strain EHA105 is described in Hood et al., *Transgen. Res.* 2: 208-218 (1993). Procedures for use of the helium gun, intact meristem preparation, tissue culture and co-cultivation conditions, and transgenic plant recovery for the intact meristem experiments are also described in Bidney et al., *supra*. Procedures for the use of the focusing gun, i.e., as in the intact meristem experiments, are described in Sautter et al., *Bio/Technol.* 9: 1080-1085 (1991). Procedures for use of the helium gun, split meristem preparation, tissue culture, co-cultivation, kanamycin selection and transgenic plant recovery for the split meristem experiments are described in Malone-Schoneberg et al., *Plant Sci.* 103: 199-207 (1994).

Parameters and general use of the focusing gun were similar to those 1.4 μm described in Sautter et al., *supra*. Gold particles measuring at a concentration of $0.5 \times 10^6 / \mu\text{l}$ and 80 bars of nitrogen were typically used although different sized particles (0.25 to 2.5 μm) and different nitrogen pressures (40-160 bar) were tested. For the Dupont PDS-1000 helium driven particle bombardment device, the meristems were bombarded twice at the highest shelf with 600 psi rupture discs and a vacuum of 26 inches Hg and 1.8 μm tungsten particles. The intact meristem transformation method involved imbibing seed for 24 hours in the dark, removing the cotyledons and root radical, followed by culturing of the meristem explants. Twenty-four hours later, the primary leaves were removed to expose the apical meristem. The explants were placed apical dome side up and bombarded twice with particles, followed by

co-cultivation with *Agrobacterium*. To start the co-cultivation for intact meristems a 0.5 μ l droplet of OD⁶⁰⁰ 4.0 *Agrobacterium* was placed on the meristem. After a 3-day co-cultivation the meristems were transferred to culture medium with 250 μ g/ml cefotaxime (+ 100 mg/l kanamycin for the NPTII selection experiments). Selection has also been done using 50 gm/l kanamycin. The split meristem method involved imbibing seed, breaking of the cotyledons to produce a clean fracture at the plane of the embryonic axis, excising the root tip and then bisecting the explants longitudinally between the primordial leaves. The two halves were placed cut surface up on the medium then bombarded twice with particles, followed by co-cultivation with *Agrobacterium*. For split meristems, after bombardment the meristems were placed in an OD⁶⁰⁰ 0.6 *Agrobacterium* suspension for 30 minutes. They were then removed from the suspension onto solid culture medium for 3-day co-cultivation. After this period, the meristems were transferred to fresh medium with 250 μ g/ml cefotaxime (+ 100mg/l kanamycin for selection). Selection has also been done using 50 gm/l kanamycin.

Table 5 summarizes the results from experiments looking at the frequency of transformed sectors in primary shoots and secondary branches. Table 6 summarizes the results from experiments carried through to progeny for analysis of seed transmission. Transformants were identified in either seed or plants by assaying for marker gene expression. For GUS expression, activity was determined using either histochemical staining or the fluorometric assay, see Jefferson et al., *EMBO J.* 6: 3901-3907 (1987). Inheritance of transgene expression was assessed using an NPTII ELISA.

TABLE 5
Overall Transformed Sector Frequencies
for Sunflower Transformation

MARKER	PLASMID PHP #	GUN	MERISTEM PREPARATION	AVE. ‡ SECTORS	REPS (N)
Screening:					
GUS	167	FG	Intact	22.0	15
GUS	167	FG	Intact	15.0	3
GUS	167	He	Intact	13.6	12
GFP	8011	He	Intact	7.5	1
Selection:					
NPTII/Kan	762	FG	Intact	12.7	3

TABLE 6
Overall Sector and Seed Transmission
for Sunflower Transformation

MARKER	PLASMID PHP #	GUN	MERISTEM PREPAR- ATION	AVE. ‡ SECTORS	REPS (N)	SEED TRANS- MISSION
Screening:						
GUS	167	FG	Intact	15.0	3	0/8
GFP	8011	He	Intact	7.5	1	9/10
Selection:						
NPTII/KAN	762	FG	Intact	12.7	4	0/2
NPTII/KAN	8011	He	Split	1.8	4	6/8

For evaluation of initial sector frequencies, GUS seemed to be the most effective marker gene (Tables 5 and 6). However, based on screening alone (no selection) the observed GUS sectors did not subsequently contribute to the germline and thus did not transmit to progeny (Table 6). It was unlikely that this was due to the use of the focusing gun because, both quantitatively and qualitatively, mapping of GUS sectors after use of each type of gun was comparable. GFP was significantly better than NPTII

(7.5% vs. 1.8%, see Table 6) for both sector frequency and seed transmission when compared using the helium gun. Progeny from GFP-expressing (no chemical selection but instead visually selecting using GFP) TO plants were analyzed for NPTII expression (Table 6) Progeny from 10 plants were tested and 9 showed inheritance of NPTII activity. From kanamycin-selected TO plants, 6 out of 8 tested showed transmission of NPTII expression to the progeny. Not only has GFP "visual selection" worked unexpectedly well for recovering the primary transformants (i.e. compare Tables 3 and 4), but cotransformation and inheritance of transgene expression was also very efficient.

Table 7: Progeny Analysis of Transformation Experiments of SMF3
with GFP (PHP8011: CaMV-GFPm, ALS-NPTII)

Treatment (Comments)	SID	No. of Seeds Screened	TO NPTII Map Range	Maximum NPTII	Transmissi on Efficiency
Topdown Non-selected (from 1 experiment, 120 explants in total)	281976	246	12.6-->30	9.9	9/10
	281977	139	16.3-->	11.9	
	281978	127	>30	>30	
	281979	175	9--27	6	
	281980	74	>30	4	
	281983	15	3--21.3	6.5	
	319454	124	4.1--18.7	5.9	
	319453	118	++	8.6	
	281982	47	>30	6.2	
	281981	3	>30	0	
Split selected (from 4 experiments, 955 explants in total)	248112	6	++	2.3	6/8
	248113	24	>30	7.8	
	248111	41	>30	>30	
	248107	23	17	2.7	
	248114	67	>30	1.2	
	248110	8	4.1	0	
	248109	6	10.7	0	
	248115	86	++	5.5	

Parallel experiments comparing GFP screening, GUS screening and NPTII/kanamycin selection can be performed using intact meristems, the Agrobacterium strain EHA105 and the helium gun, in side-by-side comparisons. Plasmid PHP8011 can be used for GFP/NPTII comparisons. An identical plasmid substituting the GUS structural gene for GFP can be used for GUS/NPTII comparisons, and both plasmids can be used in side-by-side experiments for GUS/GFP comparisons. Sector frequency in primary shoots and secondary shoots can be

assayed and calculated as described above. Inheritance and expression of transgenes can also be analyzed in these experiments.

Example 5

GFP As A FLP-Mediated Excision Marker In Maize

5 Immature embryos of the Hi-II genotype were
bombarded with PHP8674 which contains a GFPm reporter
gene that is activated upon FLP recombinase-mediated
FRT excision. Plasmid PHP8674 also carries a
10 constitutively expressed gene for bialaphos resistance.
Stably transformed callus was selected on bialaphos and
examined under UV-blue light for GFP expression. No
GFP expression was observed in these primary
transformants. Pieces of callus transformed with
15 PHP8674 were then bombarded with PHP8007. Plasmid
PHP8007 contains the FLP recombinase gene operably
linked to the ubiquitin promoter. Callus sectors
expressing GFP were first observed ten days after the
second bombardment. These GFP-expressing sectors
20 continued to grow and express the transgene after one
month of culture. As a negative control the FRT/GFPm
stably transformed callus was also bombarded with a
non-FLP containing plasmid. This treatment did not
produce GFP expressing sectors.

25 As a second control for this experiment the
consecutive treatments were done in opposite order.
Hi-II immature embryos were bombarded with PHP8007
which carries the FLP recombinase gene and a bialaphos
resistance gene. Stable transformants were recovered
following selection on medium containing bialaphos.
30 These calli were re-bombarded with PHP8674. GFP
expression was observed in the callus but this faded
over 5-10 days until it could no longer be detected.
The stably transformed callus expresses FLP
35 recombinase. Introduction of PHP8674 likely provided
the substrate for excision (the FRT sites along with
spacer DNA between the sites) which activated the GFP
cassette while still on the plasmid. This "activated

expression" was only transient and disappeared as the plasmid was degraded.

Example 6

GFP Targeting to the Nucleus as a Fusion Protein

5 Plasmid PHP9053 was constructed in which the GFP
protein was fused to the ALS enzyme (designated MALS
for maize-ALS). The ALS gene, which encodes a protein
normally found in the cytoplasm, is typically used to
confer herbicide resistance. Fang et al., *Plant Mol.*
10 *Biol.* 18: 1185-1187 (1992). A nuclear localization
signal from simian virus 40 was fused to the N terminus
of GFPm as described in Example 1 (Kalderon et al.,
Cell 39: 499-509 (1984) with the ALS enzyme fused to
the C terminus. The sequence encoding this nuclear
15 targeted fusion protein (in construct PHP9053) was
introduced into maize cells using particle bombardment
as described previously. Once gene expression was
confirmed through GFP fluorescence under the
microscope, tissue was fixed as described earlier and
20 examined to determine subcellular localization of GFP.
Fluorescence was localized to the nucleus. Plant
regeneration and assessment of toxicity for this
experiment has not yet been done.

25 GFP was targeted to the nucleus by fusing the
protein to another polypeptide that is normally
compartmentalized in the nucleus. For example, the
GFPZm-RAD51 fusion found in plasmid PHP8744 contains
RAD51, a protein required for mitotic and meiotic
recombination and double-stranded break repair, fused
30 to GFP. The gene encoding RAD51 was cloned from yeast.
Haaf et al., *Proc. Nat Acad. Sci.* 92: 2298-2302 (1995).
Degenerate primers based on the nucleotide sequence of
the RAD51 gene cloned from yeast were used to identify
and isolate clones containing the RAD51 gene from
35 maize. The carboxy terminus of GFPm was fused to the
amino terminus of ZmRAD51a. Maize cells were
transformed with the GFPm-ZmRAD51 fusion and
fluorescence was localized in the nucleus. Plant

regeneration and assessment of toxicity for this experiment has not yet been done.

Although the foregoing refers to particular preferred embodiments, it will be understood that the present invention is not so limited. It will occur to those of ordinary skill in the art that various modifications may be made to the disclosed embodiments and that such modifications are intended to be within the scope of the present invention, which is defined by the following claims.

All publications and patent applications mentioned in this specification are indicative of the level of skill of those in the art to which the invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference in its entirety.

What Is Claimed Is:

1. An isolated DNA molecule comprising a nucleotide sequence encoding a green fluorescent protein operably linked to an inducible promoter.

2. The isolated DNA sequence according to claim 1, wherein the inducible promoter is selected from the group consisting of the estrogen-inducible promoter, the estradiol-inducible promoter, the ACE1 promoter, the IN2 promoter and the tetracycline repressor promoter.

3. An isolated DNA molecule according to claim 1, wherein said nucleotide sequence encoding a green fluorescent protein is operably linked to a nucleotide sequence encoding a signal sequence for subcellular localization which directs a protein to a subcellular compartment.

4. An isolated DNA molecule comprising a nucleotide sequence selected from the group consisting of:

- (a) SEQ ID NO: 1;
- (b) a nucleotide sequence that has substantial sequence similarity with SEQ ID NO: 1; and
- (c) a functional fragment of (a) or (b), wherein said DNA molecule encodes a green fluorescent protein.

5. An isolated DNA molecule according to claim 4, wherein said nucleotide sequence encoding a green fluorescent protein is operably linked to a nucleotide sequence encoding a signal sequence for subcellular localization which directs a protein to a subcellular compartment.

6. An isolated DNA molecule according to claim 5, further comprising a promoter operably linked to said nucleotide sequence encoding a signal sequence and a green fluorescent protein.

7. An expression vector comprising said DNA molecule of claim 1.

8. An expression vector comprising said DNA molecule of claim 4.

9. The expression vector according to claim 8, wherein said nucleotide sequence encoding a green fluorescent protein is operably linked to a nucleotide sequence encoding a signal sequence which directs a protein to a subcellular compartment.

10. The expression vector according to claim 9, further comprising a first promoter operably linked to said nucleotide sequence encoding a signal sequence and a green fluorescent protein.

11. The expression vector according to claim 10, further comprising a foreign gene, wherein said foreign gene is operably linked to a second promoter.

12. A method of using said expression vector of claim 7 to produce a transformed plant, comprising the steps of introducing said DNA molecule into regenerable plant cells, inducing expression of the gene encoding the green fluorescent protein, and selecting cells containing said green fluorescent protein for regeneration.

13. A method of using said expression vector of claim 8 to produce a transformed plant, comprising the steps of introducing said expression vector into

regenerable plant cells and selecting cells containing said green fluorescent protein for regeneration.

14. The method of claim 12, wherein said regenerable plant cells are selected from the group consisting of *Zea*, *Brassica* and *Helianthus* cells.

15. The method of claim 13, wherein said regenerable plant cells are selected from the group consisting of *Zea*, *Brassica* and *Helianthus* cells.

16. A transgenic plant expressing the isolated DNA molecule of claim 1.

17. A transgenic plant comprising the vector of claim 4.

18. A method for producing a transgenic plant, comprising the steps:

- (a) constructing an expression vector comprising (i) a first promoter which is an inducible promoter, operably linked to a nucleotide sequence encoding a green fluorescent protein, and (ii) a second promoter operably linked to a foreign gene;
- (b) introducing said expression vector into regenerable plant cells;
- (c) inducing expression of the gene encoding the green fluorescent protein and selecting transformed plant cells containing said protein; and
- (d) regenerating transformed plants from said selected transformed plant cells.

19. The method of claim 18, where said inducible promoter is selected from the group consisting of the estrogen-inducible promoter, the

estradiol-inducible promoter, the ACE1 promoter, the IN2 promoter and the tetracycline repressor promoter.

20. A method for producing a transgenic plant, comprising the steps:

- (a) constructing an expression vector comprising (i) a first promoter operably linked to a nucleotide sequence encoding a sequence for subcellular localization which directs a protein to a subcellular compartment and nucleotide sequence encoding a green fluorescent protein, and (ii) a second promoter operably linked to a foreign gene;
- (b) introducing said expression vector into regenerable plant cells;
- (c) selecting transformed plant cells containing said green fluorescent protein; and
- (d) regenerating transformed plants from said selected transformed plant cells.

21. The method of claim 20, wherein the sequence for subcellular localization is selected from the group which through targeting, signaling and/or retention localize the encoded protein to mitochondria, chloroplasts, peroxisomes, vacuole, endoplasmic reticulum, cell wall or for secretion generally into the apoplast.

22. The method according to claim 18, wherein said nucleotide sequence encoding a green fluorescent protein is selected from the group consisting of:

- (a) SEQ ID NO: 1;
- (b) a nucleotide sequence that has substantial sequence similarity with SEQ ID NO: 1; and

- (c) a functional fragment of (a) or (b).

23. A method for producing a transgenic plant, comprising the steps:

- (a) constructing an expression vector comprising (i) a first promoter operably linked to a nucleotide sequence encoding a screenable marker flanked on the 5-prime and 3-prime ends with a recombinase-specific target sequence, and (ii) a second promoter operably linked to a foreign gene;
- (b) introducing said expression vector into regenerable plant cells;
- (c) selecting transformed plant cells containing said screenable marker;
- (d) transforming the plant cells with a second expression vector containing a gene encoding a site-specific recombinase;
- (e) selecting plant cells that no longer express the screenable marker;
- (f) regenerating transformed plants from said selected transformed plant cells; and
- (g) isolating said foreign protein.

24. The method according to claim 23, wherein said site-specific recombinase is selected from the group consisting of the FLP/FRT, Ac/DS and cre/lox systems.

25. The method according to claim 23, wherein said screenable marker is the green fluorescent protein.

26. An isolated DNA molecule comprising a nucleotide sequence encoding the green fluorescent

protein fused in frame to superoxide dismutase such that cells transformed with said DNA molecule contain a fusion protein that (i) produces green fluorescence in the presence of UV to blue light and (ii) displays superoxide dismutase activity.

27. A method for producing a transgenic plant comprising:

- (a) constructing an expression vector comprising (i) a first promoter operably linked to a nucleotide sequence encoding a green fluorescent protein, and (ii) a second promoter operably linked to a gene encoding an enzyme that is an oxygen scavenger, and (iii) a third promoter operably linked to a foreign gene;
- (b) introducing said expression vector into regenerable plant cells;
- (c) selecting transformed plant cells containing said green fluorescent protein; and
- (d) regenerating transformed plants from said selected transformed plant cells.

28. The method according to claim 27, wherein said oxygen scavenger enzyme is superoxide dismutase.

29. A method for producing a transgenic plant comprising:

- (a) constructing an expression vector comprising (i) a first promoter operably linked to a nucleotide sequence encoding a fusion protein comprising the green fluorescent protein and an oxygen scavenger enzyme fused in frame, and

- (ii) a second promoter operably linked to a foreign gene;
- (b) introducing said expression vector into regenerable plant cells;
- (c) selecting transformed plant cells containing said green fluorescent protein and oxygen scavenger enzyme activity; and
- (d) regenerating transformed plants from said selected transformed plant cells.


30. The method according to claim 29, wherein said oxygen scavenger enzyme is superoxide dismutase.

1/5

(Linear) MAP of: Gfpm.seq Check: 3651 from: 1 to 717

ATGTTCCAAGGGCGAGGAGCTCTTCACCGGGCTGGTGGCCATCCTCGTGGAGCTCGACGGC
1 TACAGGTTCCCGCTCCTCGAGAAGTGGCCGACCACGGGTAGGAGCACCTCGAGCTGCCG 60
M S K G E E L F T G V V P I L V E L D G
GACGTGAACGGCCACAAGTTCTCCGTGTCCGGCGAGGGCGAGGGCGACGCCACCTACGGC
61 CTGCACTTCCCGGTGTTCAAGAGGCACAGGCCGCTCCCGCTCCCGCTCGGTGGATGCCG 120
D V N G H K F S V S G E G E G D A T Y G
AAGCTCACCTCAAGTTCTCTGCCACCACGGCAAGCTCCCCGTGCCCTGGCCCCACCTC
121 TTCGAGTGGGAGTTCAAGTAGACGTGGTGGCCGTTCCGAGGGGCACGGGACCGGGTGGGAG 180
K L T L K F I C T T G K L P V P W P T L
GTGACCACCTTCTCCTACGGCGTGCAGTGTCTTCCAGGTACCCCGACCACATGAAGCAG
181 CACTGGTGAAGAGGATGCCGCACGTACGAAGAGGTCCATGGGGCTCGTGTACTTCTGTC 240
V T T F S Y G V Q C F S R Y P D H M K Q
CAGGACTTCTTCAAGTCAGCCATGCCCGAGGGCTACGTGCAGGAGAGGACCATCTTCTTC
241 GTGCTGAAGAAGTTCAAGTCGGTACGGGGCTCCCGATGCACGTCTCTCTGGTAGAAGAAG 300
H D F F K S A M P E G Y V Q E R T I F P
AAGGACGACGGCAACTACAAGACCAGCCCCGAGGTGAAGTTCCAAGGGCACACCTCGTG
301 TTCTGCTGCCGTTGATGTTCTGGTCCCGGCTCCACTTCAAGCTTCCGCTGTGGGAGCAC 360
K D D G N Y K T R A E V K F E G D T L V
AACAGGATTGAGCTCAAGGGCATCGACTTCAAGGAGGACGGCAACATCCTCGGCCACAAG
361 TTGTCCTAACTCGAGTTCCCGTAGCTGAAGTTCTCTGCGGTTGTAGGAGCCGGTGTTC 420
N R I E L K G I D F K E D G N I L G H K
CTCGAGTACAATACTCAACTCCCAACGTGTACATCATGCCCGACAAGCAGAAGAACGGC
421 GAGCTCATGTTGATGTTGAGGGTGTTCACATGTAGTACGGGCTGTTCTCTTCTTGGCC 480
L E Y N Y N S H N V Y I M A D K Q K N G
ATCAAGGTGAACCTTCAAGATCAGGCACAACATCGAGGACGGCTCAGTGCAGCTCGCTGAC
481 TAGTTCCACTTGAAGTTCTAGTCCGTGTTGTAGCTCCTGCCGAGTCAAGTCCGAGCGACTG 540
I K V N F K I R H N I E D G S V Q L A D
CACTACCAGCAGAACACCCCATCGGGCAGGGCCCCGTGCTCTCCCGACAACCACTAC
541 GTGATGGTGGTCTTGTGGGGGTAGCCGCTGCCGGGGCAGGAGCGCGCTGTTGGTGTATG 600
H Y Q Q N T P I G D G P V L L P D N H Y
CTCTCCACCCAGTCCGCCCTCTCCAAGGACCCCAAGAGAAGAGCGGACCATGGTGTCT
601 GAGAGGTGGGTCAAGCGGGAGAGGTTCTGGGGTTGCTCTTCTCCCTGGTGTACCAAGAG 660
L S T Q S A L S K D P N E K R D H M V L
CTCGAGTTCTGTACCGGTGTGGCATCACCCACGGCATGGAGGAGCTCTACAAGTGA
661 GAGCTCAAGCACTGGCGACGACCGTAGTGGGTGCCGTACCTGCTCGAGATGTTCACT 717
L E F V T A A G I T H G M D E L Y K .


FIGURE 1



PLASMIDMAP of: PHP167 Check: 6000 From: 1 To: 16750

Taglianila April 23, 1996 10:34

FIGURE 2




PHP762 (15.749 kb)

PLASMIDMAP of: PHP752.map Check: 6000 From: 1 To: 15749

Taglianila April 21, 1996 10:36


FIGURE 3



PLASMIDMAP of: Ubigfpn.Seq check: 822 from: 1 to: 5702

Taglianila April 23, 1996 10:17


FIGURE 4



PLASMIDMAP of PHP 8144. Seq check: 8061 from: 1 to: 5778

Taglianila April 23, 1996 10:22

FIGURE 5




1/5

(Linear) MAP of: Gfpa.seq Check: 1491 from: 1 to 717

[illegible]

FIGURE 1




PHP762 (15.749 kb)

FLASHMAP of: PHP762.map Check: 6000 From: 1 To: 15749

Tagliavilla April 23, 1996 10:36

FIGURE 3





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C12N 15/12, 15/82, C07K 14/435 A01H 5/00</p>	<p>A3</p>	<p>(11) International Publication Number: WO 97/41228</p> <p>(43) International Publication Date: 6 November 1997 (06.11.97)</p>
<p>(21) International Application Number: PCT/US97/07688</p> <p>(22) International Filing Date: 1 May 1997 (01.05.97)</p> <p>(30) Priority Data: 60/016,345 1 May 1996 (01.05.96) US</p> <p>(60) Parent Application or Grant (63) Related by Continuation US 60/016,345 (CIP) Filed on 1 May 1996 (01.05.96)</p> <p>(71) Applicant (for all designated States except US): PIONEER HIBRED INTERNATIONAL, INC. [US/US]; Darwin Building, 7100 N.W. 62nd Avenue, P.O. Box 1000, Johnston, IA 50131 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): GORDON-KAMM, William [US/US]; 3916 67th Street, Urbandale, IA 50322 (US). PIERCE, Dorothy, A. [US/US]; 9124 Hammontree Drive, Urbandale, IA 50322 (US). BOWEN, Benjamin [GB/US]; 3008 36th Street, Des Moines, IA 50310 (US). BIDNEY, Dennis [US/US]; 8385 Plum Drive, Urbandale, IA 50322 (US). ROSS, Margit [US/US]; 5534 Kensington</p>	<p>Circle, Johnston, IA 50131 (US). SCELONGE, Christopher [US/US]; 5314 Ovid Avenue, Des Moines, IA 50310 (US). MILLER, Mike [US/US]; 59 W. Buchanan Street, Winterset, IA 50273 (US). SANDAH, Gary [US/US]; 1649 Thornwood Road, West Des Moines, IA 50265 (US). WANG, Lijuan [CN/US]; 4728 80th Place, Urbandale, IA 50322 (US).</p> <p>(74) Agents: BENT, Stephen, A. et al.; Foley & Lardner, Suite 500, 3000 K Street, N.W., Washington, DC 20007-5109 (US).</p> <p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p> <p>(88) Date of publication of the international search report: 11 December 1997 (11.12.97)</p>	
<p>(54) Title: USE OF THE GREEN FLUORESCENT PROTEIN AS A SCREENABLE MARKER FOR PLANT TRANSFORMATION</p>		
<p>(57) Abstract</p> <p>A method for the production of transgenic plants is provided in which a vector carrying a gene encoding the green fluorescent protein is introduced into cells, the cells are screened for the protein and transformed cells are selected and regenerated. The cellular toxicity of the green fluorescent protein is circumvented by regulating expression of the gene encoding the protein or directing the protein to a subcellular compartment where it is not toxic to the cell. DNA constructs are provided for cell transformation in which the expression of a gene encoding the green fluorescent protein is placed under the control of an inducible promoter. In addition, DNA constructs are provided in which a nucleotide sequence encoding the green fluorescent protein is operably linked to a signal sequence which directs the expressed protein to a subcellular compartment where the protein is not toxic to the cell. Oxidative stress to plant cells transformed with GFP also can be ameliorated by transforming cells with an expression vector comprising genes encoding GFP and an oxygen scavenger enzyme such as superoxide dismutase. The toxicity of GFP in transformed plants can be eliminated by excising the screenable marker gene following detection of transformed cells or sectors. The FLP/FRT system is used in conjunction with GFP as a visible marker for transformation and FRT excision. A nucleotide sequence optimized for expression of the green fluorescent protein in plants is also provided.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

In International Application No

PCT/US 97/07688

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C12N15/82 C07K14/435 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHIU W.L., ET AL.: "ENGINEERED GFP AS A VITAL REPORTER IN PLANTS" CURRENT BIOLOGY, vol. 6, no. 3, pages 325-330, XP000571865 see the whole document ---	1-10, 12, 13, 16-18, 20, 22
X	SHEEN J ET AL: "GREEN-FLUORESCENT PROTEIN AS A NEW VITAL MARKER IN PLANT CELLS" PLANT JOURNAL, vol. 8, no. 5, pages 777-784, XP002003596 see the whole document ---	1, 4-10, 12, 13, 16-18, 20, 22
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

16 October 1997

Date of mailing of the international search report

31. 10. 97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Maddox, A

INTERNATIONAL SEARCH REPORT

In: tional Application No

PCT/US 97/07688

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	HAAS J., ET AL.: "Cloning vector pHGF-S65T, complete sequence, green fluorescent protein (gfp) gene, complete cds." EMBL SEQUENCE DATABASE, REL. 46, 24-JAN-1996, ACCESSION NO. U43284, XP002043798 see sequence overlap between positions 829-1545 & HAAS, J., ET AL.: "Cloning vector pHGF-S65T, complete sequence, green fluorescent protein (gfp) gene, complete cds." CURRENT BIOLOGY, vol. 6, 1996, pages 315-324, ---	4
X	HASELOFF J ET AL: "GFP IN PLANTS" TRENDS IN GENETICS, vol. 11, no. 8, page 328/329 XP002003595 see the whole document ---	4-6, 8-10,13, 20
X	HEINLEIN, M., ET AL.: "Interaction of tobamovirus movement protien with plant cytoskeleton" SCIENCE, vol. 270, 22 December 1995, pages 1983-1985, XP002043799 see the whole document ---	3
P,X	WO 96 27675 A (MEDICAL RES COUNCIL ;HASELOFF JAMES PHILLIP (AU); HODGE SARAH (GB)) 12 September 1996 see page 5, paragraph 2; example 1 see page 25, paragraph 3 - page 28, paragraph 1 see page 32, paragraph 2 ---	4-6, 8-11,13, 17,20,21
P,X	HASELOFF, J., ET AL.: "Removal of a cryptic intron and subcellular localization of green fluorescent protein are required to mark transgenic Arabidopsis brightly" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA., vol. 94, March 1997, WASHINGTON US, pages 2122-2127, XP002043800 see the whole document ---	4-6, 8-11,13, 17,20,21

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 97/07688

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	PANG, S-Z., ET AL.: "An improved green fluorescent protein gene as a vital marker in plants" PLANT PHYSIOLOGY, vol. 112, November 1996, pages 893-900, XP002043801 see the whole document	4,8,15, 17
X	YODER, J., ET AL.: "Transformation systems for generating marker-free transgenic plants" BIOTECHNOLOGY, vol. 12, no. 3, March 1994, pages 263-267, XP002043802 see the whole document	23-25
Y	---	24
Y	WO 94 03619 A (ZENECA LTD ;BRIGHT SIMON WILLIAM JONATHAN (GB); GREENLAND ANDREW J) 17 February 1994 see example 4	24
Y	WO 92 01370 A (UNIV CALIFORNIA) 6 February 1992 see page 38 - page 40	24
X	WO 93 01283 A (US) 21 January 1993 see the whole document	23,24
X	DALE, E.C., ET AL.: "Gene transfer with subsequent removal of the selection gene from the host genome" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA., vol. 88, 1991, WASHINGTON US, pages 10558-10562, XP002043803 see the whole document	23,24
X	WO 94 17176 A (PURDUE RESEARCH FOUNDATION) 4 August 1994 see the whole document	23,24
A	GOLDSBOROUGH, A.P., ET AL.: "Transposition mediated re-positioning and subsequent elimination of marker genes from transgenic tomato" BIOTECHNOLOGY, vol. 11, 1993, pages 1286-1291, XP002043843 see the whole document	23,24

	-/--	

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 97/07688

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	LLOYD, A.M., ET AL: "Functional expression of the yeast FLP/frt site-specific recombination system in <i>Nicotiana tabacum</i> ." MOL. GEN. GENET., vol. 242, 1994, pages 653-657, XP002043804 see the whole document ---	24
A	GLEAVE, A.P.: "A versatile binary vector system with a T-DNA organisational structure conducive to efficient integration of cloned DNA into the plant genome" PLANT MOLECULAR BIOLOGY, vol. 20, 1992, pages 1203-1207, XP002043805 see figure 1C ---	
E	WO 97 26333 A (UNIV FLORIDA RES FOUND ;ZOLOTUKHIN SERGEI (US); MUZYCZKA NICHOLAS) 24 July 1997 see page 8, line 17 - line 23 see page 49, line 15 - page 52, line 21 see page 62, line 25 - page 64, line 19 -----	1,2,4,5

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 97/07688

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 97 07688

FURTHER INFORMATION CONTINUED FROM PCT/ISA210

1. Claims: 1-3,7,12,14,16,18,19,22

Method for producing transgenic plants by screening for transformants on the basis of the inducible expression of green fluorecent protein. DNA constructs encoding a green fluorescent protein linked to an inducible promoter, vectors containing said constructs, and transgenic plants resulting from the method.

2. Claims: 4-6,8-11,13,15,17.

Method for producing transgenic plants by screening for transformants on the basis of expression of a plant optimized green fluorecent protein gene. Plant optimized DNA constructs encoding a green fluorescent protein, vectors containing said constructs, and transgenic plants resulting from the method.

3. Claims: 20,21.

Method for producing transgenic plants by screening for transformants on the basis of the expression of subcellular targeted green fluorecent protein.

4. Claims: 23-25

Method for producing transgenic plants by screening for transformants on the basis of the expression of green fluorecent protein, followed by sited directed excision of the marker gene using a site specific recombinase.

5. Claims: 26-30

Method for producing transgenic plants by screening for transformants on the basis of the expression of green fluorecent protein in combination with an oxygen scavenging enzyme, e.g. superoxide dismutase.

INTERNATIONAL SEARCH REPORT

In International Application No
PCT/US 97/07688

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9627675 A	12-09-96	AU 4884396 A	23-09-96
WO 9403619 A	17-02-94	AU 4718493 A	03-03-94
		EP 0658207 A	21-06-95
WO 9201370 A	06-02-92	US 5225341 A	06-07-93
		AU 660620 B	06-07-95
		AU 8102291 A	18-02-92
		EP 0577598 A	12-01-94
		JP 5508993 T	16-12-93
		US 5482852 A	09-01-96
WO 9301283 A	21-01-93	CA 2073412 A	09-01-93
WO 9417176 A	04-08-94	US 5527695 A	18-06-96
		EP 0686191 A	13-12-95
WO 9726333 A	24-07-97	NONE	

